

MODULI SPACES OF LOCAL \mathbf{G} -SHTUKAS

EVA VIEHMANN

Abstract

We give an overview of the theory of local \mathbf{G} -shtukas and their moduli spaces that were introduced in joint work of U. Hartl and the author, and in the past years studied by many people. We also discuss relations to moduli of global \mathbf{G} -shtukas, properties of their special fiber through affine Deligne-Lusztig varieties and of their generic fiber, such as the period map.

1 Introduction

Local \mathbf{G} -shtukas are an analog over local function fields of p -divisible groups with additional structure. In this article we give an overview about the theory of moduli spaces of local \mathbf{G} -shtukas and their relation to moduli of global \mathcal{G} -shtukas. It parallels the theory of Rapoport-Zink moduli spaces of p -divisible groups and their relation to Shimura varieties. Yet it has the additional charm that additional structure given by any parahoric group scheme \mathbf{G} can be easily encoded and treated in a group-theoretic way.

We begin by defining local \mathbf{G} -shtukas for a parahoric group scheme \mathbf{G} over $\text{Spec } \mathbb{F}_q[[z]]$ (see [Section 2](#)). They are pairs consisting of an $L^+\mathbf{G}$ -torsor \mathcal{G} together with an isomorphism $\sigma^*L\mathcal{G} \rightarrow L\mathcal{G}$ for the associated $L\mathbf{G}$ -torsor. Here for unexplained notation we refer to the respective sections. Local \mathbf{G} -shtukas are the function field analog of p -divisible groups with additional structure, as well as a group theoretic generalization of Drinfeld's shtukas. Next we discuss possible bounds on the singularities of local \mathbf{G} -shtukas. These are a more general replacement of the minuscule coweights fixed to define Shimura varieties. In [Section 3](#) we consider deformations for local \mathbf{G} -shtukas, and the description of the universal deformation in terms of the loop group $L\mathbf{G}$. In [Section 4](#) we describe the analog of Rapoport-Zink moduli spaces for local \mathbf{G} -shtukas bounded by a given bound \hat{Z} and within a given quasi-isogeny class. They are representable by a formal scheme, locally formally of finite type over $\text{Spf}\check{R}$. Here, \check{R} is the completion of the maximal unramified extension of the ring of definition of the chosen bound \hat{Z} . Also,

they admit a tower of coverings indexed by compact open subgroups of $\mathbf{G}(\mathbb{F}_q((z)))$. Our next topic is the special fiber of a Rapoport-Zink moduli space of local \mathbf{G} -shtukas. It can be identified with a so-called affine Deligne-Lusztig variety. In [Section 5](#) we discuss applications of this result to geometric questions such as dimensions and closure relations of Newton strata. In [Section 6](#) we describe the relation between our local \mathbf{G} -shtukas and global \mathcal{G} -shtukas where \mathcal{G} is a parahoric group scheme over a smooth projective geometrically irreducible curve C over \mathbb{F}_q . There are several results in this direction, the central idea being that one wants to describe a global \mathcal{G} -shtuka by corresponding local \mathbf{G}_i -shtukas associated with each of the legs c_i . Due to our choice of definition of local \mathbf{G} -shtukas, this works particularly well when considering global \mathcal{G} -shtukas with fixed legs. Then one obtains analogues/generalizations of Serre-Tate's theorem, as well as of several classical results comparing Shimura varieties and Rapoport-Zink spaces in the arithmetic case. In the last section we define period spaces as suitable subspaces of an affine Grassmannian. Contrary to the arithmetic case they are no longer subspaces of a classical flag variety because we allow also non-minuscule bounds. We define the period map from the analytic space associated with a Rapoport-Zink space to the corresponding period space, and discuss its image and compatibility with the tower of coverings of the generic fiber of the Rapoport-Zink space.

2 Local \mathbf{G} -shtukas

2.1 Generalities. Let \mathbb{F}_q be a finite field of characteristic p with q elements, let \mathbb{F} be a fixed algebraic closure of \mathbb{F}_q , and let $\mathbb{F}_q[[z]]$ and $\mathbb{F}_q[[\zeta]]$ be the power series rings over \mathbb{F}_q in the (independent) variables z and ζ . As base schemes we will consider the category $\mathrm{Nilp}_{\mathbb{F}_q[[\zeta]]}$ consisting of schemes over $\mathrm{Spec} \mathbb{F}_q[[\zeta]]$ on which ζ is locally nilpotent. Let \mathbf{G} be a parahoric group scheme over $\mathrm{Spec} \mathbb{F}_q[[z]]$ with connected reductive generic fiber, compare [Bruhat and Tits \[1972\]](#), Déf. 5.2.6 and [Haines and Rapoport \[2008\]](#).

Let $S \in \mathrm{Nilp}_{\mathbb{F}_q[[\zeta]]}$ and consider any sheaf of groups H on S for the fpqc-topology. By an H -torsor on S we mean a sheaf \mathcal{H} for the fpqc-topology on S together with a (right) action of the sheaf H such that \mathcal{H} is isomorphic to H on an fpqc-covering of S .

Let now $L\mathbf{G}$ and $L^+\mathbf{G}$ be the loop group and the group of positive loops associated with \mathbf{G} , i.e. for an \mathbb{F}_q -algebra R let

$$(L^+\mathbf{G})(R) = \mathbf{G}(R[[z]]) \quad \text{and} \quad (L\mathbf{G})(R) = \mathbf{G}(R((z))).$$

Let \mathcal{G} be an $L^+\mathbf{G}$ -torsor on S . Via the inclusion $L^+\mathbf{G} \subset L\mathbf{G}$ we can associate an $L\mathbf{G}$ -torsor $L\mathcal{G}$ with \mathcal{G} . For any $L\mathbf{G}$ -torsor \mathcal{G}' on S we denote by $\sigma^*\mathcal{G}'$ the pullback of \mathcal{G}' under the q -Frobenius morphism $\sigma := \mathrm{Frob}_q : S \rightarrow S$.

Definition 2-1. A *local \mathbf{G} -shtuka* over some $S \in \text{Nilp}_{\mathbb{F}_q}[[\xi]]$ is a pair $\underline{\mathcal{G}} = (\mathcal{G}, \tau_{\mathcal{G}})$ consisting of an $L^+\mathbf{G}$ -torsor \mathcal{G} on S and an isomorphism of the associated $L\mathbf{G}$ -torsors $\tau_{\mathcal{G}} : \sigma^*L\mathcal{G} \xrightarrow{\sim} L\mathcal{G}$.

A *quasi-isogeny* $g : (\mathcal{G}', \tau_{\mathcal{G}'}) \rightarrow (\mathcal{G}, \tau_{\mathcal{G}})$ between local \mathbf{G} -shtukas over S is an isomorphism $g : L\mathcal{G}' \xrightarrow{\sim} L\mathcal{G}$ of the associated $L\mathbf{G}$ -torsors with $g \circ \tau_{\mathcal{G}'} = \tau_{\mathcal{G}} \circ \sigma^*g$.

Local \mathbf{G} -shtukas were introduced and studied in [Hartl and Viehmann \[2011\]](#), [Hartl and Viehmann \[2012\]](#) in the case where \mathbf{G} is a constant split reductive group over \mathbb{F}_q . The general case was first considered in work of [Arasteh Rad and Hartl \[2014\]](#).

Example 2-2. For $\mathbf{G} = \text{GL}_r$, we have the following more classical description.

A local shtuka over $S \in \text{Nilp}_{\mathbb{F}_q}[[\xi]]$ (of rank r) is a pair (M, ϕ) where M is a sheaf of $\mathcal{O}_S[[z]]$ -modules on S which Zariski-locally is free of rank r , together with an isomorphism of $\mathcal{O}_S((z))$ -modules

$$\phi : \sigma^*M \otimes_{\mathcal{O}_S[[z]]} \mathcal{O}_S((z)) \xrightarrow{\sim} M \otimes_{\mathcal{O}_S[[z]]} \mathcal{O}_S((z)).$$

Then the category of local GL_r -shtukas over S is equivalent to the category of local shtukas of rank r over S , see [Hartl and Viehmann \[2011\]](#), Lemma 4.2.

Local shtukas were first introduced by [Anderson \[1993\]](#) over a complete discrete valuation ring. [Genestier \[1996\]](#) constructed moduli spaces for them in the Drinfeld case and used these to uniformize Drinfeld modular varieties.

An important invariant of local \mathbf{G} -shtukas is their *Newton point*. To define it let k be an algebraically closed field of characteristic p and let $L = k((z))$. Then every local \mathbf{G} -shtuka $(\mathcal{G}, \tau_{\mathcal{G}})$ over k has a trivialization $\mathcal{G} \cong (L^+\mathbf{G})_k$. Via this isomorphism the Frobenius map $\tau_{\mathcal{G}}$ corresponds to an element $b \in L\mathbf{G}(k) = \mathbf{G}(L)$. Changing the trivialization replaces b by a different representative of its $L^+\mathbf{G}(k)$ - σ -conjugacy class. In the same way, local \mathbf{G} -shtukas isogenous to $(\mathcal{G}, \tau_{\mathcal{G}})$ correspond to the elements of the σ -conjugacy class

$$[b] = \{g^{-1}b\sigma(g) \mid g \in \mathbf{G}(L)\}.$$

Let G denote the generic fiber of \mathbf{G} . Then the set of σ -conjugacy classes $B(G) = \{[b] \mid b \in G(L)\}$ for quasi-split G is described by [Kottwitz \[1985\]](#), [Kottwitz \[1997\]](#) by two invariants. Let T be a maximal torus of G . The first invariant is then the Kottwitz point, i.e. the image under the Kottwitz map $\kappa_G : G(L) \rightarrow \pi_1(G)_{\Gamma}$. Here, $\pi_1(G)$ is Borovoi's fundamental group, and Γ is the absolute Galois group of $\mathbb{F}_q((z))$. This invariant is locally constant on $L\mathbf{G}$, and can also be computed using the identification $\pi_0(L\mathbf{G}) = \pi_1(G)/\Gamma$, compare [Neupert \[2016\]](#), 2.2. The second invariant is the Newton point ν_b , an element of $X_*(T)_{\mathbb{Q}}^{\Gamma}$.

Consider now a local \mathbf{G} -shtuka $(\mathcal{G}, \tau_{\mathcal{G}})$ over a scheme S . Then we obtain an induced decomposition of $S(k)$ into subsets $\mathfrak{N}_{[b]}(k)$ for $[b] \in B(G)$ with

$$\mathfrak{N}_{[b]}(k) = \{x \in S(k) \mid (\mathcal{G}, \tau_{\mathcal{G}})_x \text{ is in the isogeny class of } [b]\}.$$

By [Rapoport and Richartz \[1996\]](#), this induces a decomposition of S into locally closed subschemes, which we call Newton strata. If S is connected, they correspond to the strata obtained by fixing the Newton point only.

2.2 Bounds. To ensure finiteness properties of moduli spaces of local \mathbf{G} -shtukas, or of deformations of local \mathbf{G} -shtukas, we bound the singularity of the morphism $\tau_{\mathcal{G}}$. In the arithmetic context, such bounds are usually assumed to be minuscule and correspond to the choice of the Hodge cocharacter defining a Shimura datum. In our context we define bounds more generally as suitable ind-subschemas of the affine flag variety.

The *affine flag variety* $\text{Flag}_{\mathbf{G}}$ associated with G is the fpqc-sheaf associated with the presheaf

$$S \mapsto L\mathbf{G}(S)/L^+\mathbf{G}(S) = \mathbf{G}(\mathcal{O}_S((z))(S))/\mathbf{G}(\mathcal{O}_S[[z]](S))$$

on the category of \mathbb{F}_q -schemes. By [Pappas and Rapoport \[2008\]](#), Theorem 1.4 and [Richarz \[2016\]](#) Theorem A, $\text{Flag}_{\mathbf{G}}$ is represented by an ind-scheme which is ind-projective over \mathbb{F}_q .

Consider further the group scheme $\mathbf{G} \times_{\mathbb{F}_q[[z]]} \text{Spec } \mathbb{F}_q((\zeta))[[z - \zeta]]$ under the homomorphism

$$(2-3) \quad \mathbb{F}_q[[z]] \rightarrow \mathbb{F}_q((\zeta))[[z - \zeta]], \quad z \mapsto z = \zeta + (z - \zeta).$$

The associated affine Grassmannian $\text{Gr}_{\mathbf{G}}^{\text{dR}}$ is the sheaf of sets for the fpqc-topology on $\text{Spec } \mathbb{F}_q((\zeta))$ associated with the presheaf

$$(2-4) \quad X \mapsto \mathbf{G}(\mathcal{O}_X((z - \zeta)))/\mathbf{G}(\mathcal{O}_X[[z - \zeta]]).$$

Remark 2-5. $\text{Gr}_{\mathbf{G}}^{\text{dR}}$ is in the same way as $\text{Flag}_{\mathbf{G}}$ above an ind-projective ind-scheme over $\text{Spec } \mathbb{F}_q((\zeta))$. Here, note that the homomorphism in (2-3) induces an inclusion $\mathbb{F}_q((z)) \rightarrow \mathbb{F}_q((\zeta))[[z - \zeta]]$, so the group $\mathbf{G} \times_{\mathbb{F}_q[[z]]} \text{Spec } \mathbb{F}_q((\zeta))[[z - \zeta]]$ is reductive, justifying the name affine ‘‘Grassmannian’’. The notation B_{dR} refers to the fact that if C is the completion of an algebraic closure of $\mathbb{F}_q((\zeta))$, then $C((z - \zeta))$ is the function field analog of Fontaine’s p -adic period field B_{dR} .

We fix an algebraic closure $\overline{\mathbb{F}_q((\zeta))}$ of $\mathbb{F}_q((\zeta))$, and consider pairs (R, \hat{Z}_R) , where $R/\mathbb{F}_q[[\zeta]]$ is a finite extension of discrete valuation rings contained in $\overline{\mathbb{F}_q((\zeta))}$, and where $\hat{Z}_R \subset \widehat{\text{Flag}}_{\mathbf{G},R} := \text{Flag}_{\mathbf{G}} \widehat{\otimes}_{\mathbb{F}_q} \text{Spf } R$ is a closed ind-subscheme that is contained in a

bounded (or: projective) subscheme of $\widehat{\text{Flag}}_{\mathbf{G},R}$. Two such pairs (R, \hat{Z}_R) and $(R', \hat{Z}'_{R'})$ are equivalent if they agree over a finite extension of the rings R, R' .

A *bound* is then abstractly defined as an equivalence class $\hat{Z} := [(R, \hat{Z}_R)]$ of such pairs (R, \hat{Z}_R) satisfying certain properties (for the precise conditions compare [Hartl and Viehmann \[2017\]](#), Def. 2.1), in particular

1. $\hat{Z}_R \subset \widehat{\text{Flag}}_{\mathbf{G},R}$ is a ζ -adic formal scheme over $\text{Spf}R$ that is stable under the left $L^+\mathbf{G}$ -action.
2. The special fiber $Z_R := \hat{Z}_R \widehat{\times}_{\text{Spf}R} \text{Spec } \kappa_R$ is a quasi-compact subscheme of $\text{Flag}_G \widehat{\times}_{\mathbb{F}_q} \kappa_R$ where κ_R is the residue field of R .
3. Let \hat{Z}_R^{an} be the strictly $R[\frac{1}{\zeta}]$ -analytic space associated with \hat{Z}_R . Then the \hat{Z}_R^{an} are invariant under the left multiplication of $\mathbf{G}(\cdot[[z - \zeta]])$ on $\text{Gr}_{\mathbf{G}}^{\text{BdR}}$.

The reflex ring $R_{\hat{Z}}$ of $[(R, \hat{Z}_R)]$ is the intersection in $\overline{\mathbb{F}_q((\zeta))}$ of the fixed field of $\{\gamma \in \text{Aut}_{\mathbb{F}_q[[\zeta]]}(\overline{\mathbb{F}_q((\zeta))}) \mid \gamma(\hat{Z}) = \hat{Z}\}$ with all finite extensions of $\mathbb{F}_q[[\zeta]]$ over which a representative \hat{Z}_R of the given bound exists.

Let $(\mathfrak{g}, \tau_{\mathfrak{g}})$ be a local \mathbf{G} -shtuka over some $S \in \text{Nilp}_{R_{\hat{Z}}}$ and let S' be an étale covering of S over which a trivialization $\mathfrak{g} \cong (L^+\mathbf{G})_{S'}$ exists. Then $(\mathfrak{g}, \tau_{\mathfrak{g}})$ (or $\tau_{\mathfrak{g}}$) is *bounded by* \hat{Z} if for every such trivialization and for every finite extension R of $\mathbb{F}_q[[\zeta]]$ over which a representative \hat{Z}_R of \hat{Z} exists, the morphism

$$S' \widehat{\times}_{R_{\hat{Z}}} \text{Spf}R \rightarrow LG \widehat{\times}_{\mathbb{F}_q} \text{Spf}R \rightarrow \widehat{\text{Flag}}_{\mathbf{G},R}$$

induced by $\tau_{\mathfrak{g}}$ factors through \hat{Z}_R .

The above definition of bounds follows the strategy to allow as general bounds as possible. On the other hand, classically and by analogy with the arithmetic case, one considers bounds given as Schubert varieties. They are described as follows (compare [Arasteh Rad and Hartl \[2014\]](#), Example 4.12).

Example 2-6. Consider the base change G_L of G to $L = \mathbb{F}((z))$. Let A be a maximal split torus in G_L and let T be its centralizer, a maximal torus. Let $N = N(T)$ be the normalizer of T and let \mathcal{T}^0 be the identity component of the Néron model of T over $\mathcal{O}_L = \mathbb{F}[[z]]$.

Consider the Iwahori–Weyl group $\widetilde{W} = N(L)/\mathcal{T}^0(\mathcal{O}_L)$ and let $\widetilde{W}^{\mathbf{G}} = (N(L) \cap \mathbf{G}(\mathcal{O}_L))/\mathcal{T}^0(\mathcal{O}_L)$. Then by the Bruhat–Tits decomposition we have a bijection

$$(2-7) \quad \widetilde{W}^{\mathbf{G}} \backslash \widetilde{W} / \widetilde{W}^{\mathbf{G}} \rightarrow L^+\mathbf{G}(\mathbb{F}) \backslash LG(\mathbb{F}) / L^+\mathbf{G}(\mathbb{F}).$$

Let ω be in the right hand side, and let \mathbb{F}_{ω} be a finite extension of \mathbb{F}_q such that ω has a representative $g_{\omega} \in LG(\mathbb{F}_{\omega})$. We define the *Schubert variety* S_{ω} as the ind-scheme theoretic closure of the $L^+\mathbf{G}$ -orbit of g_{ω} in $\text{Flag}_{\mathbf{G}} \widehat{\times}_{\mathbb{F}_q} \mathbb{F}_{\omega}$. Let $R = \mathbb{F}_{\omega}[[\zeta]]$ and let $\hat{Z}_{\mathbb{F}_{\omega}[[\zeta]]} =$

$S_\omega \widehat{\times}_{\mathbb{F}_\omega} \mathrm{Spf} R$. Then the equivalence class of (R, \hat{Z}_R) defines a bound. In this case we also say “bounded by ω ” instead of “bounded by (R, \hat{Z}_R) ”.

3 Deformations

Let $\mathrm{Flag}_{\mathbf{G}}$ denote again the affine flag variety, i.e. the quotient sheaf $\mathrm{Flag}_{\mathbf{G}} = L\mathbf{G}/L^+\mathbf{G}$, an ind-scheme over \mathbb{F}_q which is of ind-finite type. Let $\widehat{\mathrm{Flag}}_{\mathbf{G}}$ be the fiber product $\mathrm{Flag}_{\mathbf{G}} \times_{\mathrm{Spec} \mathbb{F}_q} \mathrm{Spf} \mathbb{F}_q((\xi))$.

Then generalizing [Hartl and Viehmann \[2011, Prop. 5.1\]](#) (where the case that \mathbf{G} is split is considered) the ind-scheme $\widehat{\mathrm{Flag}}_{\mathbf{G}}$ pro-represents the functor $\mathrm{Nilp}_{\mathbb{F}_q[[\xi]]}^o \rightarrow (\mathrm{Sets})$

$$S \mapsto \{(\mathcal{G}, \delta) \mid \mathcal{G} \text{ a } \mathbf{G}\text{-torsor on } S, \\ \delta : L\mathcal{G} \rightarrow L\mathbf{G}_S \text{ an isomorphism of the associated } L\mathbf{G}\text{-torsors}\} / \cong .$$

Here (\mathcal{G}, δ) and (\mathcal{G}', δ') are isomorphic if $\delta^{-1} \circ \delta'$ is an isomorphism $\mathcal{G}' \rightarrow \mathcal{G}$.

We fix a bound $[(R, \hat{Z}_R)]$ and a local \mathbf{G} -shtuka $\underline{\mathbb{G}} = (\mathbb{G}, \phi_{\mathbb{G}})$ over a field $k \in \mathrm{Nilp}_{\mathbb{F}_q[[\xi]]}$ which is bounded by \hat{Z}_R . We consider the functor of bounded deformations of $\underline{\mathbb{G}}$ on the category of Artinian local $k[[\xi]]$ -algebras with residue field k ,

$$\mathrm{Def}_{\underline{\mathbb{G}}, \hat{Z}_R} : (\mathrm{Art}_k) \rightarrow (\mathrm{Sets}) \\ A \mapsto \{(\underline{\mathcal{G}}, \beta) \mid \underline{\mathcal{G}} \text{ a local } \mathbf{G}\text{-shtuka over } \mathrm{Spec} A \text{ bounded by } \hat{Z}_R \\ \beta : \underline{\mathbb{G}} \rightarrow \underline{\mathcal{G}} \otimes_A k \text{ an isomorphism of local } \mathbf{G}\text{-shtukas}\} / \cong$$

and require isomorphisms to be compatible with the maps β .

Remark 3-1. In [Hartl and Viehmann \[ibid.\]](#), 5 this functor and the deformation space representing it are explicitly described in case that \mathbf{G} is split and \hat{Z}_R associated with some $\omega \in \tilde{W}$ as in [Example 2-6](#), see below. This is generalized to reductive groups \mathbf{G} in [Viehmann and Wu \[2016\]](#). It would be interesting to have these results also in the more general context of parahoric group schemes.

Assume that \mathbf{G} is a reductive group scheme over $\mathrm{Spec} \mathbb{F}_q[[z]]$. Then the left hand side of [Equation \(2-7\)](#) is $W_0 \backslash \tilde{W} / W_0$ where W_0 is the finite Weyl group of $\mathbf{G}_{\mathbb{F}_q}$. This double quotient is isomorphic to $X_*(T)_{\mathrm{dom}}$. Let $\mu \in X_*(T)_{\mathrm{dom}}$ and (R, \hat{Z}_R) be the corresponding bound as in [Example 2-6](#). Let $\underline{\mathbb{G}}$ be a local \mathbf{G} -shtuka bounded by μ over a field $k \in \mathrm{Nilp}_{\mathbb{F}_q[[\xi]]}$. Assume that there is a trivialization $\alpha : \underline{\mathbb{G}} \rightarrow (\mathbf{G}_k, b_0\sigma)$ for some $b_0 \in L\mathbf{G}(k)$. Using the boundedness, let $x \in Z_R(k)$ be the point defined by b_0^{-1} . By [Hartl and Viehmann \[2011\]](#), Thm. 5.6 and [Viehmann and Wu \[2016\]](#), Proposition 2.6 we have

Theorem 3-2. *Let \mathbf{G} be reductive, and \hat{Z}_R as in [Example 2-6](#). Let \mathbb{G} and α be as above. Let D be the complete local ring of \hat{Z}_R at the point x . It is a complete noetherian local ring over $k[[\zeta]]$. Then D pro-represents the formal deformation functor $\text{Def}_{\mathbb{G}, \hat{Z}_R}$.*

To study Newton strata in a family of local \mathbf{G} -shtukas it is helpful to also have a corresponding stratification on deformations. However, for infinitesimal deformations such a question does not make sense. Instead we will replace the formal deformation space $\text{Spf} D$ by $\text{Spec } D$, using the following proposition. Here, a linearly topologized $\mathbb{F}_q[[z]]$ -algebra R is admissible if $R = \varprojlim R_\alpha$ for a projective system $(R_\alpha, u_{\alpha\beta})$ of discrete rings such that the filtered index-poset has a smallest element 0, all maps $R \rightarrow R_\alpha$ are surjective, and the kernels $I_\alpha := \ker u_{\alpha,0} \subset R_\alpha$ are nilpotent.

Proposition 3-3. *Let R be an admissible $\mathbb{F}_q[[z]]$ -algebra as above with filtered index-poset \mathbb{N}_0 . Then the pullback under the natural morphism $\text{Spf} R \rightarrow \text{Spec } R$ defines a bijection between local \mathbf{G} -shtukas bounded by μ over $\text{Spec } R$ and over $\text{Spf} R$.*

Remark 3-4. Without a boundedness condition the pullback map is in general only injective. The corresponding result for p -divisible groups is shown by Messing and by de Jong. The above proposition is proved for split \mathbf{G} in [Hartl and Viehmann \[2011\]](#), Proposition 3.16. However, the same proof also shows this assertion for all \mathbf{G} that are reductive over $\text{Spec } \mathbb{F}_q[[z]]$.

4 Moduli spaces

In this section we fix \mathbf{G} , an isogeny class of local \mathbf{G} -shtukas and a bound \hat{Z} . We then define moduli spaces of local \mathbf{G} -shtukas bounded by \hat{Z} in the same way as Rapoport-Zink define their moduli spaces of p -divisible groups in [Rapoport and Zink \[1996\]](#).

Definition 4-1. Let \mathbb{G}_0 be a local \mathbf{G} -shtuka over \mathbb{F} . Let $\hat{Z} = [(R, \hat{Z}_R)]$ be a bound and denote its reflex ring by $R_{\hat{Z}} = \kappa[[\xi]]$. It is a finite extension of $\mathbb{F}_q[[\xi]]$. Set $\check{R}_{\hat{Z}} := \mathbb{F}[[\xi]]$, and consider the functor

$$\begin{aligned} \check{\mathfrak{M}} : (\text{Nilp}_{\check{R}_{\hat{Z}}})^\circ &\rightarrow (\text{Sets}) \\ S &\mapsto \{(\underline{\mathfrak{g}}, \bar{\delta}) \mid \underline{\mathfrak{g}} \text{ a local } \mathbf{G}\text{-shtuka over } S \text{ bounded by } \hat{Z}^{-1}, \\ &\quad \bar{\delta} : \underline{\mathfrak{g}}_{\bar{S}} \rightarrow \mathbb{G}_{0, \bar{S}} \text{ a quasi-isogeny}\} / \cong \end{aligned}$$

Here $\bar{S} := V_S(\zeta)$ is the zero locus of ζ in S , and two pairs $(\underline{\mathfrak{g}}, \bar{\delta}), (\underline{\mathfrak{g}}', \bar{\delta}')$ are isomorphic if $\bar{\delta}^{-1} \circ \bar{\delta}'$ lifts to an isomorphism $\underline{\mathfrak{g}}' \rightarrow \underline{\mathfrak{g}}$.

By [Hartl and Viehmann \[2017\]](#), 2.2 $\underline{\mathfrak{g}}$ is bounded by \hat{Z}^{-1} if and only if $\tau_{\mathfrak{g}}^{-1}$ is bounded by \hat{Z}_R .

Let $\text{QIsog}_{\mathbb{F}}(\underline{\mathbb{G}}_0)$ be the group of self-quasi-isogenies of $\underline{\mathbb{G}}_0$. It naturally acts on the functor $\check{\mathfrak{M}}$. Since \mathbb{F} has no non-trivial étale coverings, we may fix a trivialization $\underline{\mathbb{G}}_0 \cong ((L^+\mathbf{G})_{\mathbb{F}}, b\sigma^*)$ where $b \in L\mathbf{G}(\mathbb{F})$ represents the Frobenius morphism. Via such a trivialization, $\text{QIsog}_{\mathbb{F}}(\underline{\mathbb{G}}_0)$ is identified with

$$(4-2) \quad J_b(\mathbb{F}_q((z))) := \{g \in G(\mathbb{F}((z))) \mid g^{-1}b\sigma(g) = b\}.$$

This is the set of $\mathbb{F}_q((z))$ -valued points of an algebraic group J_b over $\mathbb{F}_q((z))$.

Theorem 4-3 (Arasteh Rad and Hartl [2014], Thm. 4.18 and Cor. 4.26, Hartl and Viehmann [2017], Rem. 3.5). *The functor $\check{\mathfrak{M}}$ is ind-representable by a formal scheme over $\text{Spf}\check{R}_{\check{Z}}$ which is locally formally of finite type and separated. It is an ind-closed ind-subscheme of $\text{Flag}_{\mathbf{G}} \widehat{\times}_{\mathbb{F}_q} \text{Spf}\check{R}_{\check{Z}}$.*

By its analogy with moduli spaces of p -divisible groups, the formal scheme representing $\check{\mathfrak{M}}$ is called a Rapoport-Zink space for bounded local \mathbf{G} -shtukas.

Let E be the quotient field of $R_{\check{Z}}$, and $\check{E} \cong \mathbb{F}((\xi))$ the completion of its maximal unramified extension. We write $\check{\mathfrak{M}}^{\text{an}}$ for the strictly \check{E} -analytic space associated with $\check{\mathfrak{M}}$.

Next we explain the construction of a tower of coverings of $\check{\mathfrak{M}}^{\text{an}}$. Roughly spoken, it is obtained by trivializing the (dual) Tate module of the universal local \mathbf{G} -shtuka over $\check{\mathfrak{M}}^{\text{an}}$.

Definition 4-4. Let S be an $\mathbb{F}((\zeta))$ -scheme or a strictly $\mathbb{F}((\zeta))$ -analytic space. Then an étale local \mathbf{G} -shtuka over S is a pair $\underline{\mathfrak{g}} = (\mathfrak{g}, \tau_{\mathfrak{g}})$ consisting of an $L^+\mathbf{G}$ -torsor \mathfrak{g} on S and an isomorphism $\tau_{\mathfrak{g}} : \sigma^*\mathfrak{g} \rightarrow \mathfrak{g}$ of $L^+\mathbf{G}$ -torsors.

Remark 4-5. Let $S = \text{Spf}B$ be an affinoid admissible formal $\check{R}_{\check{Z}}$ -scheme and S^{an} the associated strictly \check{E} -analytic space. Consider a trivialized local \mathbf{G} -shtuka $((L^+\mathbf{G})_S, A\sigma^*)$ over S . Then $A \in \mathbf{G}(B[[z]][\frac{1}{z-\zeta}])$. Note that

$$(z - \zeta)^{-1} = - \sum_{i=0}^{\infty} \zeta^{-i-1} z^i \in \mathcal{O}_{S^{\text{an}}}(S^{\text{an}})[[z]]$$

implies $B[[z]][\frac{1}{z-\zeta}] \subset \mathcal{O}_{S^{\text{an}}}(S^{\text{an}})[[z]]$. Therefore $((L^+\mathbf{G})_S, A\sigma^*)$ induces an étale local \mathbf{G} -shtuka $((L^+\mathbf{G})_{S^{\text{an}}}, A\sigma^*)$ over S^{an} .

To obtain similarly a universal family of étale local \mathbf{G} -shtukas over $\check{\mathfrak{M}}^{\text{an}}$, we cover $\check{\mathfrak{M}}$ by affinoid admissible formal $\check{R}_{\check{Z}}$ -schemes. For each of them one chooses a finite étale covering trivializing the local \mathbf{G} -shtuka, and applies the above construction. Descending the étale local \mathbf{G} -shtuka then yields the desired universal family, compare Hartl and Viehmann [ibid.], 6.

There are two approaches to construct the (dual) Tate module of an étale local \mathbf{G} -shtuka $\underline{\mathfrak{g}}$ over a connected strictly \check{E} -analytic space X . To define it directly, following Neupert

[2016], 2.6, consider for each $n \in \mathbb{N}$ the $L^+\mathbf{G}/\mathbf{G}_n$ -torsor associated with \mathfrak{G} where \mathbf{G}_n is the kernel of the projection $\mathbf{G}(\mathbb{F}_q[[z]]) \rightarrow \mathbf{G}(\mathbb{F}_q[[z]]/(z^n))$. The isomorphisms $\tau_{\mathfrak{G}} : \sigma^*\mathfrak{G} \rightarrow \mathfrak{G}$ and $\sigma^* : \mathfrak{G} \rightarrow \sigma^*\mathfrak{G}$ then induce corresponding maps of $L^+\mathbf{G}/\mathbf{G}_n$ -torsors. The invariants of $\tau \circ \sigma^*$ form a $\mathbf{G}(\mathbb{F}_q[[z]]/(z^n))$ -torsor which is trivialized by a finite étale covering of X . One can then define the Tate module of $\underline{\mathfrak{G}}$ as the inverse limit over n of these torsors.

Alternatively, one can define it as a tensor functor, following [Arasteh Rad and Hartl \[2014\]](#), or [Hartl and Viehmann \[2017\]](#), 7: Fix a geometric base point \bar{x} of X . We consider representations $\rho : \mathbf{G} \rightarrow \mathrm{GL}_r$ in $\mathrm{Rep}_{\mathbb{F}_q[[z]]}(\mathbf{G})$. Let $\underline{M} = (M, \tau_M)$ be the étale local shtuka of rank r associated with $\rho_*\underline{\mathfrak{G}}$ as in [Example 2-2](#). Let $\underline{M}_{\bar{x}}$ denote its fiber over \bar{x} . The (dual) Tate module of $\underline{\mathfrak{G}}$ with respect to ρ is the (dual) Tate module of $\underline{M}_{\bar{x}}$,

$$\check{T}\underline{M}_{\bar{x}} = \{m \in \underline{M}_{\bar{x}} \mid \tau_M(\sigma^*m) = m\},$$

a free $\mathbb{F}_q[[z]]$ -module of rank r . It carries a continuous monodromy action of $\pi_1^{\mathrm{ét}}(X, \bar{x})$, which also factors through $\pi_1^{\mathrm{alg}}(X, \bar{x})$. We obtain the *dual Tate module* of $\underline{\mathfrak{G}}$ as a tensor functor

$$\check{T}_{\underline{\mathfrak{G}}, \bar{x}} : \mathrm{Rep}_{\mathbb{F}_q[[z]]}\mathbf{G} \rightarrow \mathrm{Rep}_{\mathbb{F}_q[[z]]}^{\mathrm{cont}}(\pi_1^{\mathrm{alg}}(X, \bar{x})).$$

Similarly, using rational representations in $\mathrm{Rep}_{\mathbb{F}_q((z))}\mathbf{G}$ one can define the *rational dual Tate module*

$$\check{V}_{\underline{\mathfrak{G}}, \bar{x}} : \mathrm{Rep}_{\mathbb{F}_q((z))}\mathbf{G} \rightarrow \mathrm{Rep}_{\mathbb{F}_q((z))}^{\mathrm{cont}}(\pi_1^{\mathrm{ét}}(X, \bar{x})).$$

The two constructions of dual Tate modules are compatible in the sense that the tensor functor associated with the above torsor coincides with the tensor functor $\check{T}_{\underline{\mathfrak{G}}, \bar{x}} : \mathrm{Rep}_{\mathbb{F}_q[[z]]}\mathbf{G} \rightarrow (\mathbb{F}_q[[z]]\text{-Loc})_X$ with values in the local systems of $\mathbb{F}_q[[z]]$ -lattices on X that can be associated with $\check{T}_{\underline{\mathfrak{G}}, \bar{x}}$ as in the second construction (compare [Hartl and Viehmann \[ibid.\]](#), Prop. 5.3).

We can now proceed to define level structures and the tower of coverings of $\check{\mathfrak{M}}^{\mathrm{an}}$. Let FMod_A denote the category of finite locally free A -modules. We consider the forgetful functors

$$(4-6) \quad \omega_A^\circ : \mathrm{Rep}_A\mathbf{G} \rightarrow \mathrm{FMod}_A$$

and

$$(4-7) \quad \mathit{forget} : \mathrm{Rep}_A^{\mathrm{cont}}(\pi_1^{\mathrm{ét}}(X, \bar{x})) \rightarrow \mathrm{FMod}_A.$$

For an étale local \mathbf{G} -shtuka $\underline{\mathfrak{G}}$ over X the two sets

$$\begin{aligned} \mathrm{Triv}_{\underline{\mathfrak{G}}, \bar{x}}(\mathbb{F}_q[[z]]) &= \mathrm{Isom}^{\otimes}(\omega_{\mathbb{F}_q[[z]]}^\circ, \mathit{forget} \circ \check{T}_{\underline{\mathfrak{G}}, \bar{x}})(\mathbb{F}_q[[z]]) \\ \mathrm{Triv}_{\underline{\mathfrak{G}}, \bar{x}}(\mathbb{F}_q((z))) &= \mathrm{Isom}^{\otimes}(\omega_{\mathbb{F}_q((z))}^\circ, \mathit{forget} \circ \check{V}_{\underline{\mathfrak{G}}, \bar{x}})(\mathbb{F}_q((z))) \end{aligned}$$

are non-empty and carry natural actions of $\mathbf{G}(\mathbb{F}_q[[z]]) \times \pi_1^{\text{alg}}(X, \bar{x})$ and $G(\mathbb{F}_q((z))) \times \pi_1^{\text{ét}}(X, \bar{x})$, respectively. Here, the first factor acts via ω° and the second via the action on the Tate module. Note that non-emptiness needs our assumption that \mathbf{G} has connected fibers, and thus slightly differs from the setting of Rapoport and Zink.

In the same way as in the arithmetic case we define coverings of $\check{\mathfrak{M}}$.

Definition 4-8. Let $\underline{\mathfrak{g}}$ be an étale local \mathbf{G} -shtuka over a connected \check{E} -analytic space X , and let K be an open compact subgroup of $\mathbf{G}(\mathbb{F}_q[[z]])$. An *integral K -level structure* on $\underline{\mathfrak{g}}$ is a $\pi_1^{\text{alg}}(X, \bar{x})$ -invariant K -orbit in $\text{Triv}_{\underline{\mathfrak{g}}, \bar{x}}(\mathbb{F}_q[[z]])$.

For an open subgroup $K \subset \mathbf{G}(\mathbb{F}_q[[z]])$ let X^K be the functor on the category of \check{E} -analytic spaces over X parametrizing integral K -level structures on the local \mathbf{G} -shtuka $\underline{\mathfrak{g}}$ over X .

Let $K \subset \mathbf{G}(\mathbb{F}_q((z)))$ be compact open. Let $K' \subset K$ be a normal subgroup of finite index with $K' \subset \mathbf{G}(\mathbb{F}_q[[z]])$. Then $gK' \in K/K'$ acts on $X^{K'}$ by sending the triple $(\underline{\mathfrak{g}}, \delta, \eta K')$ over $X^{K'}$ to the triple $(\underline{\mathfrak{g}}, \delta, \eta gK')$.

We define $\check{\mathfrak{M}}^K$ as the \check{E} -analytic space which is the quotient of $\check{\mathfrak{M}}^{K'} := \check{\mathfrak{M}}^{\text{an}, K'}$ by the finite group K/K' . It is independent of the choice of K' . In particular, $\check{\mathfrak{M}}^{K_0} = (\check{\mathfrak{M}})^{\text{an}}$ for $K_0 = \mathbf{G}(\mathbb{F}_q[[z]])$.

By Hartl and Viehmann [2017], Cor. 7.13 one can also describe $\check{\mathfrak{M}}^K$ directly as a parameter space of local \mathbf{G} -shtukas with level structure.

Furthermore, if $K' \subseteq K$ is as above, the action of $gK' \in K/K'$ on $\check{\mathfrak{M}}^{K'}$ can be described in this interpretation as a Hecke correspondence.

5 The geometry of the special fiber

We fix again a bound (R, \hat{Z}_R) and let κ be the residue field of R . Let Z^{-1} be the special fiber of the inverse bound \hat{Z}^{-1} over κ . The affine Deligne-Lusztig variety associated with an element $b \in L\mathbf{G}(\mathbb{F})$ and Z^{-1} is the reduced closed ind-subscheme $X_{Z^{-1}}(b) \subset \text{Flag}_{\mathbf{G}}$ whose k -valued points (for any field extension k of \mathbb{F}) are given by

$$(5-1) \quad X_{Z^{-1}}(b)(k) = \{g \in \text{Flag}_{\mathbf{G}}(k) \mid g^{-1}b\sigma(g) \in Z^{-1}(k)\}.$$

Our conditions on the bound imply that Z^{-1} is a left- $L^+\mathbf{G}$ -invariant subscheme of $\text{Flag}_{\mathbf{G}}$. The $L^+\mathbf{G}$ -orbits in $\text{Flag}_{\mathbf{G}}$ correspond bijectively to the elements of $\widetilde{W}^{\mathbf{G}} \setminus \widetilde{W} / \widetilde{W}^{\mathbf{G}}$, compare Example 2-6. For $x \in \widetilde{W}$ let

$$(5-2) \quad X_x(b)(k) = \{g \in \text{Flag}_{\mathbf{G}}(k) \mid g^{-1}b\sigma(g) \in L^+\mathbf{G}xL^+\mathbf{G}\}.$$

Thus by left invariance every $X_{Z^{-1}}(b)(k)$ is a union of affine Deligne-Lusztig varieties of the form $X_x(b)$. By the boundedness condition on \hat{Z}_R , this union is finite. In the

arithmetic context, such unions of affine Deligne-Lusztig varieties have been studied for example in [He \[2016\]](#).

Affine Deligne-Lusztig varieties are the underlying reduced subschemes of Rapoport-Zink spaces:

Theorem 5-3 ([Hartl and Viehmann \[2011\]](#), 6, [Arasteh Rad and Hartl \[2014\]](#), Thm. 4.18, Cor. 4.26). *The underlying reduced subscheme of the moduli space $\check{\mathfrak{M}}$ associated with \hat{Z}_R and b as in [Theorem 4-3](#) equals $X_{Z^{-1}}(b)$. It is a scheme locally of finite type and separated over \mathbb{F} , all of whose irreducible components are projective.*

Thus to describe the special fiber of the Rapoport-Zink spaces $\check{\mathfrak{M}}$ it is enough to study the affine Deligne-Lusztig varieties $X_x(b)$. For an overview of the current state of the art of this field compare [Viehmann \[2015\]](#) and X. He’s talk at this ICM [He \[2018\]](#). We will here only discuss one class of results that were obtained as an application of the relation to local \mathbf{G} -shtukas, and that were one of the initial goals of this theory. For the following result it is essential to assume that \mathbf{G} is reductive over $\mathbb{F}_q[[z]]$. For more general groups there are counterexamples to all assertions in the theorem. Recall that for reductive groups $\widetilde{W}^{\mathbf{G}} \setminus \widetilde{W} / \widetilde{W}^{\mathbf{G}} \cong X_*(T)_{\text{dom}}$. Let $\text{def}(b) = \text{rk } G - \text{rk}_{\mathbb{F}_q((z))} J_b$ where J_b is as in (4-2). Note that $X_*(T)_{\text{dom}}$ is partially ordered by the Bruhat ordering. Via the induced ordering on the Newton points (requiring equality of Kottwitz points) also $B(G)$ inherits an ordering. Let l denote the length of a maximal chain between two elements in the partially ordered set $B(G)$, and let ρ be the half-sum of the positive roots of \mathbf{G} .

Theorem 5-4 ([Viehmann \[2013\]](#), [Hamacher and Viehmann \[2017\]](#), 3.2). *Let $\mu_1 \preceq \mu_2 \in X_*(T)$ be dominant coweights and $[b] \in B(G)$. Assume that $\kappa_G(b) = \mu_2$ in $\pi_1(G)_{\Gamma}$ and that $v_b \preceq \mu_2$.*

1. Let

$$S_{\mu_1, \mu_2} = \bigcup_{\mu_1 \preceq \mu' \preceq \mu_2} L^+ \mathbf{G} \mu'(z) L^+ \mathbf{G}.$$

Then the Newton stratum $\mathfrak{N}_{[b]} = [b] \cap S_{\mu_1, \mu_2}$ is pure of codimension

$$l([b], [\mu_2]) = \langle \rho, \mu_2 - v_b \rangle + \frac{1}{2} \text{def}(b)$$

in S_{μ_1, μ_2} . The closure of $\mathfrak{N}_{[b]}$ in S_{μ_1, μ_2} is the union of all $\mathfrak{N}_{[b']}$ for $[b']$ with $\kappa_G(b') = \mu_2$ in $\pi_1(\mathbf{G})_{\Gamma}$ and $v_{b'} \preceq v_b$.

2. $X_{\mu_2}(b)$ and $X_{\preceq \mu_2}(b) = \bigcup_{\mu \preceq \mu_2} X_{\mu}(b)$ are equidimensional of dimension

$$\dim X_{\mu_2}(b) = \dim X_{\preceq \mu_2}(b) = \langle \rho, \mu_2 - v_b \rangle - \frac{1}{2} \text{def}(b).$$

Here, the first assertion is most useful for $\mu_1 = \mu_2$, or for μ_1 the unique minuscule coweight with $\mu_1 \preceq \mu_2$.

Note that one needs to define the notions of codimension and of the closure of the infinite-dimensional schemes $\mathfrak{N}_{[b]}$. Both of these definitions use that there is an open subgroup H of LG^+ such that the Newton point of an element of S_{μ_1, μ_2} only depends on its H -coset, an element of the finite-dimensional scheme $S_{\mu_1, \mu_2}/H$, compare [Viehmann \[2013\]](#), 4.3. These (co)dimensions are directly linked to the dimensions of Newton strata in the universal deformation of a local \mathbf{G} -shtuka as in [Section 3](#). In fact, [Theorem 5-4](#) is equivalent to a completely analogous result on dimensions and closure relations in that context, and this equivalence is also used in the proof of [Theorem 5-4](#).

The proofs of the two parts of the theorem are closely linked: One first shows the upper bound on $\dim X_{\preceq \mu_2}(b)$ and how to compute $\dim \mathfrak{N}_{[b]}$ from $\dim X_{\preceq \mu_2}(b)$. Then a purity result for the Newton stratification proves the lower bound and equidimensionality. For the proof compare [Hamacher and Viehmann \[2017\]](#), 3.2 explaining how the proof for split \mathbf{G} in [Viehmann \[2013\]](#) can be generalized. The theorem also inspired a corresponding theory for Newton strata in Shimura varieties of Hodge type by Hamacher.

6 Comparison to global \mathcal{G} -shtukas

6.1 Global \mathcal{G} -shtukas. Moduli spaces of bounded global \mathcal{G} -shtukas are the function field analogue of Shimura varieties. Using these moduli spaces [Lafforgue \[2012\]](#) showed one direction of the global Langlands correspondence for all groups \mathcal{G} in the function field case. In this section we describe the relation between moduli spaces of bounded local \mathbf{G} -shtukas and global \mathcal{G} -shtukas.

To define them, let C be a smooth projective geometrically irreducible curve over \mathbb{F}_q , and let \mathcal{G} be a parahoric group scheme over C , i.e. a smooth affine group scheme with connected fibers whose generic fiber is reductive over $\mathbb{F}_q(C)$ and such that for every point v of C for which the fiber above v is not reductive, the group scheme \mathcal{G}_v is a parahoric group scheme over A_v , as defined by Bruhat and Tits. Here A_v denotes the completion of $k[C]$ at v .

Definition 6-1. A *global shtuka* with n legs over a scheme S is a tuple $(\mathcal{G}, s_1, \dots, s_n, \varphi)$ where

1. $\mathcal{G} \in \mathcal{H}^1(C, \mathcal{G})(S)$ is a \mathcal{G} -torsor over $C \times_{\mathbb{F}_q} S$
2. the $s_i \in C(S)$ are pairwise disjoint S -valued points, called the legs
3. φ is an isomorphism

$$\varphi : \sigma^* \mathcal{G}|_{C \times_{\mathbb{F}_q} S \setminus \{\cup_i \Gamma_{s_i}\}} \rightarrow \mathcal{G}|_{C \times_{\mathbb{F}_q} S \setminus \{\cup_i \Gamma_{s_i}\}},$$

called the Frobenius isomorphism.

The stack $\nabla_n \mathcal{H}^1(C, \mathcal{G})$ over $\text{Spec } \mathbb{F}_q$ is the *moduli stack of global \mathcal{G} -shtukas with n legs*.

There is a canonical morphism $\nabla_n \mathcal{H}^1(C, \mathcal{G}) \rightarrow C^n \setminus \Delta$ mapping a global \mathcal{G} -shtuka to its legs.

We fix distinct places $c_i \in C(\mathbb{F})$ for $1 \leq i \leq n$, and write $c = (c_1, \dots, c_n)$. Let A_c be the completion of the local ring $\mathcal{O}_{C^n, c}$, and let \mathbb{F}_c be the residue field. Then $A_c \cong \mathbb{F}_c \llbracket \zeta_1, \dots, \zeta_n \rrbracket$.

Writing $A_{c_i} \cong \mathbb{F}_{c_i} \llbracket z \rrbracket$, let $\mathbf{G}_{c_i} = \mathcal{G} \times_C \text{Spec } A_{c_i}$, a parahoric group scheme over $\text{Spec } \mathbb{F}_{c_i} \llbracket z \rrbracket$, and $\mathbf{G}_i := \text{Res}_{\mathbb{F}_{c_i} \llbracket z \rrbracket / \mathbb{F}_q \llbracket z \rrbracket} \mathbf{G}_{c_i}$.

Let

$$\nabla_n \mathcal{H}^1(C, \mathcal{G})^c = \nabla_n \mathcal{H}^1(C, \mathcal{G}) \widehat{\times}_{C^n} \text{Spf} A_c$$

be the formal completion of the stack $\nabla_n \mathcal{H}^1(C, \mathcal{G})$ along c . It parametrizes global \mathcal{G} -shtukas with n legs in the formal neighborhoods of the c_i . We want to define a global-local functor associating with such a global \mathcal{G} -shtuka local shtukas at each of these places, compare [Arasteh Rad and Hartl \[2014\]](#), or [Neupert \[2016\]](#), 3.2. Let $(\mathcal{G}, (s_i), \varphi) \in \nabla_n \mathcal{H}^1(C, \mathcal{G})^c(S)$. In other words, $s_i : S \rightarrow C$ factors through $\text{Spf} A_{c_i}$. We consider each place c_i separately. Let $S_{c_i} = \text{Spec } \mathbb{F}_{c_i} \times_{\text{Spec } \mathbb{F}_q} S$. The base change of \mathcal{G} to $\text{Spf} A_{c_i} \widehat{\times}_{\mathbb{F}_q} S = S_{c_i} \llbracket z \rrbracket$ defines an element of $\mathcal{H}^1(\mathbb{F}_q, \mathbf{G}_{c_i})(S_{c_i} \llbracket z \rrbracket)$. We now use the following translation, cf. [Hartl and Viehmann \[2011\]](#), Prop. 2.2a, [Neupert \[2016\]](#), Prop. 3.2.4, or [Arasteh Rad and Hartl \[2014\]](#), 2.4.

Proposition 6-2. *Let S be a scheme over \mathbb{F}_q . There is a natural equivalence of categories*

$$\mathcal{H}^1(\mathbb{F}_q, \mathbf{G})(S \llbracket z \rrbracket) \rightarrow \mathcal{H}^1(\mathbb{F}_q, L^+ \mathbf{G})(S).$$

We thus obtain an element of

$$\mathcal{H}^1(\mathbb{F}_q, L^+ \mathbf{G}_{c_i})(S_{c_i}) = \mathcal{H}^1(\mathbb{F}_q, L^+ \mathbf{G}_i)(S),$$

or an $L^+ \mathbf{G}_i$ -torsor \mathcal{G}_i over S .

Also, the Frobenius morphism φ induces local Frobenius morphisms φ_i on the \mathbf{G}_i -torsors for each i . Altogether we obtain a functor

$$\mathfrak{L} = (\mathfrak{L}_1, \dots, \mathfrak{L}_n) : \nabla_n \mathcal{H}^1(C, \mathcal{G})^c \rightarrow \prod_i \text{Sht}_{\mathbf{G}_i}, \quad (\mathcal{G}, (s_i), \varphi) \mapsto ((\mathcal{G}_i, \varphi_i))$$

called the global-local functor.

In the same way, one can associate with every global \mathcal{G} -shtuka $\underline{\mathcal{G}} \in \nabla_n \mathcal{H}^1(C, \mathcal{G})^c(S)$ for some scheme S and every fixed place $s \in C \setminus \{c_i\}$ an étale local \mathbf{G}_s -shtuka $\mathfrak{L}_s(\underline{\mathcal{G}}) =$

$\underline{\mathcal{G}}_S$. One can then define level structures on global \mathcal{G} -shtukas away from the legs using the corresponding notion for local \mathbf{G} -shtukas. For a detailed discussion compare [Neupert \[2016\]](#), 3.4.

Definition 6-3. Let $\underline{\mathcal{G}} = (\mathcal{G}, (s_i), \varphi)$ be a global \mathcal{G} -shtuka over S , let $D_0 \subset C$ be a finite reduced subscheme with $s_i \in (C \setminus D_0)(S)$ for every i . For every $v \in D_0$ fix an open subgroup $U_v \subseteq L^+G_v(\mathbb{F}_q)$ and let $U = \prod U_v$. Then an *integral U -level structure* on $\underline{\mathcal{G}}$ consists of an integral U_v -level structure of $\mathcal{L}_v(\underline{\mathcal{G}})$ for every $v \in D_0$.

We denote by $\nabla \mathcal{H}_U^1(C, \mathcal{G})^c$ the *stack of global \mathcal{G} -shtukas* in $\nabla \mathcal{H}^1(C, \mathcal{G})^c$ with U -level structure.

Rather ad hoc boundedness conditions are defined as follows (compare [Arasteh Rad and Hartl \[2013\]](#)).

Definition 6-4. Let $\hat{Z}_c = (\hat{Z}_i)_i$ be a tuple of bounds $\hat{Z}_i \subset \widehat{\text{Flag}}_{\mathbf{G}_i}$. Then $(\mathcal{G}, (s_i), \varphi) \in \nabla_n \mathcal{H}^1(C, \mathcal{G})^c(S)$ is *bounded by \hat{Z}_c* if for every i the associated local \mathbf{G}_i -shtuka $(\mathcal{G}_i, \varphi_i)$ is bounded by \hat{Z}_i . We denote by $\nabla_n^{\hat{Z}_c} \mathcal{H}^1(C, \mathcal{G})^c$ the substack of $\nabla_n \mathcal{H}^1(C, \mathcal{G})^c$ of global \mathcal{G} -shtukas bounded by \hat{Z}_c , and analogously for $\nabla_n^{\hat{Z}_c} \mathcal{H}_U^1(C, \mathcal{G})^c$.

There are two other definitions of boundedness: A (seemingly more natural) one by [Varshavsky \[2004\]](#), Def. 2.4b for which, however, it is not clear how to establish a compatibility between bounded global and local shtukas. In [Neupert \[2016\]](#), 3.3 Neupert gives a global definition of boundedness, generalizing that of Varshavsky and he shows that it coincides with the one presented here.

6.2 The Serre-Tate theorem. Classically, the Serre-Tate theorem gives an equivalence of categories between deformations of an abelian variety over a scheme on which p is locally nilpotent, and its p -divisible group. By work of Arasteh Rad and Hartl, the analog in our situation also holds, taking into account that shtukas have several legs, and requiring them to be in the formal neighborhoods of fixed places of C .

Let S be in Nilp_{A_c} and let $j : \bar{S} \rightarrow S$ be a closed subscheme defined by a locally nilpotent sheaf of ideals. Let $c_1, \dots, c_n \in C$ and let $(\mathcal{G}, (s_i), \varphi) \in \nabla_n \mathcal{H}^1(C, \mathcal{G})^c(\bar{S})$. Let $((\mathcal{G}_i, \varphi_i)) = \mathcal{L}(\mathcal{G}, (s_i), \varphi)$.

Let $\text{Def}_S((\mathcal{G}, (s_i), \varphi))$ be the category of lifts of $(\mathcal{G}, (s_i), \varphi)$ to S , i.e. of pairs (\mathcal{H}, α) where $\mathcal{H} \in \nabla_n \mathcal{H}^1(C, \mathcal{G})^c(S)$ and where α is an isomorphism between $(\mathcal{G}, (s_i), \varphi)$ and the base change of \mathcal{H} to \bar{S} . Morphisms of lifts are isomorphisms compatible with the morphisms α . For the local \mathbf{G}_i -shtukas we define analogously a category of lifts $\text{Def}_S((\mathcal{G}_i, \varphi_i))$.

Theorem 6-5 ([Arasteh Rad and Hartl \[2014\]](#), Th. 5.10). *Let $(c_i)_i \in C^n$, let $(\mathcal{G}, (s_i), \varphi) \in \nabla_n \mathcal{H}^1(C, \mathcal{G})^c(\bar{S})$ and let $((\mathcal{G}_i, \varphi_i)) = \mathcal{L}(\mathcal{G}, (s_i), \varphi)$. Then the global-local functor induces an equivalence of categories $\text{Def}_S((\mathcal{G}, (s_i), \varphi)) \rightarrow \prod_i \text{Def}_S((\mathcal{G}_i, \varphi_i))$.*

6.3 Foliations. We begin by considering the uniformization morphism for Newton strata by Rapoport-Zink spaces, following [Neupert \[2016\]](#), 5 and [Arasteh Rad and Hartl \[2014\]](#), 5, and paralleling the uniformization result for p -divisible groups of [Rapoport and Zink \[1996\]](#). In this subsection we assume that \mathcal{G} is the base change to C of a reductive group G over \mathbb{F}_q . It would be interesting to have a generalization of this result to the more general context considered before.

Let T denote a maximal torus of G , and fix a Borel subgroup B containing T . Let C be as above and fix characteristic places $c_1, \dots, c_n \in C$. Let $\mu_i \in X_*(T)$ be dominant cocharacters defined over a finite extension E of \mathbb{F}_q , and $\mu = (\mu_1, \dots, \mu_n) \in X_*(T)^n$. Fix decent local \mathbf{G}_i -shtukas $\underline{\mathbf{G}}_i = (L^+ \mathbf{G}_{iE}, b_i \sigma^*)$ over $\text{Spec } E$, in particular $b_i \in L\mathbf{G}_i(E)$. Let $\mathfrak{M}_{b_i}^{\leq \mu_i}$ be the Rapoport-Zink space associated with $[b_i]$ and the bound given by μ_i .

Theorem 6-6 ([Neupert \[2016\]](#), Theorem 5.1.18, [Arasteh Rad and Hartl \[2013\]](#), Theorem 7.4). *Let S be a DM-stack over $\text{Spf}E[[\zeta_1, \dots, \zeta_n]]$ such that $(\zeta_1, \dots, \zeta_n)$ is locally nilpotent on S . Let $(\mathcal{G}_0, \phi_0, \psi_0) \in \nabla_n \mathcal{H}_U^1(C, \mathcal{G})^c(S)$ for some congruence subgroup U . For each place c_i assume that there is, and fix, an isomorphism $\mathcal{L}_{c_i}(\mathcal{G}_0, \phi_0) \cong \underline{\mathbf{G}}_i$. Then there is a morphism of formal DM-stacks over $\text{Spf}E[[\zeta_1, \dots, \zeta_n]]$*

$$S \times_{\text{Spf}E[[\zeta_1, \dots, \zeta_n]]} \prod_i \mathfrak{M}_{b_i}^{\leq \mu_i} \rightarrow \nabla_n^\mu \mathcal{H}_U^1(C, \mathcal{G})^c$$

which is ind-proper and formally étale.

This theorem has also generalizations to coverings of the moduli spaces associated with compatible level structures, and is then equivariant with respect to the action of $\mathcal{G}(\mathbb{A}_{\mathbb{Q}}^{c_i})$ by Hecke correspondences.

We consider for the universal global \mathcal{G} -shtuka $\underline{\mathcal{G}}$ over $\nabla_n^\mu \mathcal{H}_U^1(C, \mathcal{G})^c$ the strata

$$\mathfrak{N}_{\underline{\mathcal{G}}}^{([b_i])_i} = \bigcap_i \mathfrak{N}_{[b_i], \mathcal{L}_i(\underline{\mathcal{G}})} \subseteq \nabla_n^\mu \mathcal{H}_U^1(C, \mathcal{G})^c.$$

Here, the $\mathfrak{N}_{[b_i], \mathcal{L}_i(\underline{\mathcal{G}})}$ are the Newton strata in $\nabla_n^\mu \mathcal{H}_U^1(C, \mathcal{G})^c$ associated with the local \mathbf{G}_i -shtuka $\mathcal{L}_i(\underline{\mathcal{G}})$.

Then the uniformization morphism of [Theorem 6-6](#) maps each geometric point of the left hand side to $\mathfrak{N}_{\underline{\mathcal{G}}}^{([b_i])_i}$ where the $[b_i]$ are the classes of the elements used to define the Rapoport-Zink space. Refining this description we now consider a foliation structure on these Newton strata, following [Neupert \[2016\]](#), 5.

In the arithmetic case, Oort and later [Mantovan \[2004\]](#) defined a foliation structure on Newton strata on Shimura varieties, by Rapoport-Zink spaces and so-called Igusa varieties. The latter are defined as covers of a central leaf, i.e. the locus in the Shimura variety

where the p -divisible group is in a very particular isomorphism class. In our situation, the definition is slightly more involved. One needs to pass to the perfection of all involved moduli spaces to define the analog of Igusa varieties, and then to construct the foliation morphism.

Recall that a Newton stratum only depends on $L\mathbf{G}_i$ - σ -conjugacy classes $[b_i]$, and not on individual representatives. A fundamental alcove in $[b_i]$ is an element x_{b_i} of \tilde{W} such that all (or one) representative $b_{i,0}$ in $N(L)$ is contained in $[b_i]$, and such that the length $\ell(x_{b_i})$ is minimal with that property, compare Viehmann [2014], Theorem 6.5, or Nie [2015]. One can then show that this length is equal to $\langle 2\rho_G, \nu_{b_i} \rangle$ where ρ_G is the half-sum of the positive roots of G and where ν_{b_i} is the dominant Newton point of $[b_i]$. Fundamental alcoves are a group-theoretic generalization of the minimal p -divisible groups studied by Oort. For each $[b_i]$ we fix such a representative $b_{i,0}$. The central leaf

$$\mathfrak{C}_U^{(b_{i,0})} \subseteq \mathfrak{n}_{\underline{g}}^{([b_i])_i}$$

is then defined as the locus where the associated local \mathbf{G}_i -shtukas are isomorphic to $((L^+\mathbf{G}_i)_k, b_{i,0})$. It depends on the choice of the fundamental alcove, but not on that of the representative in $N(L)$.

We now define formal versions of these subschemes of $\nabla_n^\mu \mathcal{H}_U^1(C, G)$.

Definition 6-7. Recall that the formal completion of $C^n \setminus (c_1, \dots, c_n)$ is isomorphic to $\mathrm{Spf} A_c = \mathrm{Spf} \mathbb{F}_c \llbracket \zeta_1, \dots, \zeta_n \rrbracket$.

Let $\mathfrak{N}_U^{([b_i])_i}$ denote the formal completion of $\nabla_n^\mu \mathcal{H}_U^1(C, G)^c$ along the Newton stratum $\mathfrak{n}_{\underline{g}}^{([b_i])_i}$. This is not $(\zeta_1, \dots, \zeta_n)$ -adic.

Let $\mathfrak{C}_U^{(b_{i,0})}$ be the locus in $\nabla_n^\mu \mathcal{H}_U^1(C, G)^c$ such that after an fpqc-covering each local \mathbf{G}_i -shtuka associated with the universal global \mathcal{G} -shtuka is isomorphic to $(L^+\mathbf{G}_i, b_{i,0}\sigma^*)$. One can then show that this is represented by a $(\zeta_1, \dots, \zeta_n)$ -adic formal scheme over $\mathrm{Spf} A_c$, called the *central leaf*. (Neupert [2016], Prop. 6.1.7)

By X^\sharp we denote the perfection of a formal scheme X . It is again a formal scheme.

For $d \in \mathbb{N}$ we consider the subgroups $K_d = \{g \in L^+\mathbf{G}_i \mid g \equiv 1 \pmod{z^{d+1}}\}$ and

$$I_d(b_{i,0}) = \bigcap_{N \geq 0} \phi^N(K_d)$$

where $\phi(g) = b_{i,0}^{-1}\sigma^{-1}(g)b_{i,0}$.

The following theorem shows existence of the formal Igusa varieties.

Theorem 6-8 (Neupert [ibid.], Cor. 6.1.9). *For all tuples $(d_i)_i$, define $\mathfrak{I}_{\mathbf{G}_U}^{(d_i)\sharp}(T)$ as the set of pairs consisting of an element of $\mathfrak{C}_U^{(b_i)\sharp}(T)$ together with I_{d_i} -truncated isomorphisms between the associated local \mathbf{G}_i -shtukas and $(L^+\mathbf{G}_i, b_{i,0}\sigma^*)$ for all i . Then this*

defines a sheaf $\mathfrak{S}_{\mathfrak{G}_U}^{(d_i)\#}$ over $\mathfrak{C}_U^{(b_i)\#}$ representable by a $(\zeta_1, \dots, \zeta_n)$ -adic formal scheme which is finite étale over $\mathfrak{C}_U^{(b_i)\#}$. It is called a formal Igusa variety of level (d_i) .

Similarly, $\mathfrak{S}_{\mathfrak{G}_U}^{(\infty_i)\#}$ parametrizing trivializations of the whole local \mathbf{G}_i -shtukas is representable by a formal scheme, isomorphic to $\varprojlim_{\leftarrow} \mathfrak{S}_{\mathfrak{G}_U}^{(d_i)\#}$.

The foliation structure is then established by the following result.

Theorem 6-9 (Neupert [ibid.], Thm. 6.2.1). *There is a natural formally étale morphism of formal schemes over $\mathrm{Spf}A_c^\#$*

$$\hat{\pi}_{(\infty_i)} : \prod_i \mathfrak{M}_{b_{i,0}}^{\leq \mu_i \#} \times_{\mathrm{Spf}A_c^\#} \mathfrak{S}_{\mathfrak{G}_U}^{(\infty_i)\#} \rightarrow \mathfrak{M}_U^{[b_i]\#}.$$

There are also variants of this result for the special fiber of the spaces, using bounds on the quasi-isogenies, and for the associated adic spaces in the sense of Huber.

The very rough idea of the construction of this morphism is the following: A point in the Igusa variety gives us a global shtuka, such that at each place c_i the associated local \mathbf{G}_i -shtuka is the one defined by $b_{i,0}$. Then a point in the Rapoport-Zink space $\mathfrak{M}_{b_{i,0}}^{\leq \mu_i}$ gives us a modification by a quasi-isogeny of this local \mathbf{G}_i -shtuka, and we define a new global \mathfrak{G} -shtuka by keeping the old one away from the c_i and replacing it in a formal neighborhood of c_i by this new local \mathbf{G}_i -shtuka.

Applying the theorem to the cohomology of the moduli space of global \mathfrak{G} -shtukas, and its decomposition into Newton strata, one can express the cohomology of Newton strata, and therefore of the whole moduli space of global \mathfrak{G} -shtukas in terms of the cohomology of Rapoport-Zink moduli spaces and of Igusa varieties. For technical reasons the statement is only known over a finite extension E' of \mathbb{F}_q that can be described in terms of \mathfrak{G} and μ . Denote by K the function field of the curve C , by \overline{K} an algebraic closure, and by $\Gamma_{E'}$ the absolute Galois group of E' .

Theorem 6-10 (Neupert [ibid.], Main Theorem 2). *Let $\nabla_n^\mu \mathcal{H}^1(C, \mathfrak{G})$ be a moduli space of global \mathfrak{G} -shtukas, such that all connected components of $\nabla_n^\mu \mathcal{H}^1(C, \mathfrak{G}) \times_{C^n \setminus \Delta} \mathrm{Spec} E'[\zeta_1, \dots, \zeta_n]$ are proper over $\mathrm{Spec} E'[\zeta_1, \dots, \zeta_n]$.*

Then there exists a canonical isomorphism between the virtual $\mathfrak{G}(\mathbb{A}_{\mathbb{Q}}^{c_i}) \times \Gamma_{E'}$ -representations

$$\sum_j (-1)^j H_c^j(\nabla_n^\mu \mathcal{H}^1(C, \mathfrak{G}) \times \overline{K}, \mathbb{Q}_\ell)$$

and

$$\sum_{[b_i]} \sum_{d,e,f} (-1)^{d+e+f} \text{Tor}_d^{\mathfrak{K}(\prod J_{b_i,0})} \left(H_c^e \left(\prod \mathfrak{M}_{b_i,0}^{\leq \mu_i} \times \mathbb{F}, R\Psi_\eta^{\text{an}} \mathbb{Q}_\ell \right), \lim_U \lim_{d_i} H_c^f \left(\text{Ig}_U^{(d_i)} \times \mathbb{F}, R\Psi_\eta^{\text{an}} \mathbb{Q}_\ell \right) \right).$$

7 Period spaces and the period map

In our context, period spaces are constructed as strictly $\mathbb{F}_q((\zeta))$ -analytic spaces in the sense of Berkovich. In the generality presented here, they are introduced and studied in Hartl and Viehmann [2017]. We allow more general bounds than those associated with minuscule coweights. Even if the bound is given by a coweight as in Example 2-6, the bound is in general a union of Schubert cells, and not a single one as in the minuscule case. For these two reasons the period spaces have to be defined as subspaces of an affine Grassmannian instead of a (classical) flag variety. To define them, we consider the affine Grassmannian Gr_G^{Bdr} of Remark 2-5. The space of Hodge-Pink \mathbf{G} -structures bounded by \hat{Z} is defined as $\mathfrak{H}_{\mathbf{G},\hat{Z}} := \hat{Z}_E$. It is a projective subscheme of $\text{Gr}_G^{\text{Bdr}} \widehat{\times}_{\mathbb{F}_q((\zeta))} E$. Fix a local \mathbf{G} -shtuka \underline{G}_0 over \mathbb{F} and a trivialization $\underline{G}_0 \cong (L^+ \mathbf{G}_{\mathbb{F}}, b\sigma^*)$.

A z -isocrystal over \mathbb{F} is a pair (D, τ_D) consisting of a finite-dimensional $\mathbb{F}((z))$ -vector space D and an isomorphism $\tau_D : \sigma^* D \rightarrow D$ of $\mathbb{F}((z))$ -vector spaces. A Hodge-Pink structure on (D, τ_D) over a field extension L of $\mathbb{F}((\zeta))$ is a free $L[[z - \zeta]]$ -submodule $\mathfrak{q}_D \subset D \otimes_{\mathbb{F}((z))} L((z - \zeta))$ of full rank, compare Hartl and Kim [2016], 5.

Definition 7-1. Let L and b be as above and let $\gamma \in \text{Gr}_G^{\text{Bdr}}(L)$. Let $\rho : G_{\mathbb{F}_q((z))} \rightarrow \text{GL}_{n,\mathbb{F}_q((z))}$ be a representation and let V be the representation space. Then we associate with b (and ρ) the z -isocrystal $(V \otimes_{\mathbb{F}_q((z))} \mathbb{F}((z)), \rho(\sigma^* b)\sigma^*)$ over \mathbb{F} . With γ we associate the Hodge-Pink structure over L defined by

$$\mathfrak{q}_D(V) = \rho(\gamma) \cdot V \otimes_{\mathbb{F}_q((z))} L[[z - \zeta]] \subset D \otimes_{\mathbb{F}((z))} L((z - \zeta)).$$

Let

$$\underline{D}_{b,\gamma}(V) = (V \otimes_{\mathbb{F}_q((z))} \mathbb{F}((z)), \rho(\sigma^* b)\sigma^*, \mathfrak{q}_D(V)).$$

Consider the Newton point $v_b \in X_*(T)$ as a homomorphism $\mathbb{D}_{\mathbb{F}((z))} \rightarrow G_{\mathbb{F}((z))}$. Recall the Kottwitz map $\kappa_G : G(\mathbb{F}((z))) \rightarrow \pi_1(G)_\Gamma$. We then define the Hodge degree and the Newton degree as

$$t_H(\underline{D}_{b,\gamma}(V)) = \rho_*(\kappa_G(\gamma)) \quad \text{and} \quad t_N(\underline{D}_{b,\gamma}(V)) = \det_V \circ \rho \circ v_b.$$

Using $\pi_1(\mathrm{GL}(V))_\Gamma = \mathbb{Z}$ and $\mathrm{Hom}(\mathbb{D}_{\mathbb{F}((z))}, \mathbf{G}_m) = \mathbb{Q}$ we can view these as rational numbers. (Compare [Hartl and Viehmann \[2017\]](#), 4 for a definition in terms of $\underline{D}_{b,\gamma}(V)$ only.)

We call $\underline{D}_{b,\gamma}(V)$ *weakly admissible* if the images of $[b]$ and γ in $\pi_1(G)_{\Gamma, \mathbb{Q}}$ coincide and for all subobjects the Hodge degree is less or equal to the Newton degree. Finally, the pair (b, γ) is weakly admissible if $\underline{D}_{b,\gamma}(V)$ is weakly admissible for a faithful representation ρ of G .

Let $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}} := \check{\mathcal{H}}_{\mathbf{G}, \hat{Z}} \widehat{\times}_E \check{E}$ where $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}} = \hat{Z}_E$. The *period space* $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^{wa}$ is then defined as the set of all γ in the associated \check{E} -analytic space $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}}^{\mathrm{an}}$ such that (b, γ) is weakly admissible.

To define the admissible locus $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^a \subset \check{\mathcal{H}}_{\mathbf{G}, \hat{Z}}^{\mathrm{an}}$, one associates with \underline{D} a pair of σ -bundles $\underline{\mathfrak{E}}(\underline{D}) = \underline{\mathfrak{E}}_{b,\gamma}(V)$ and $\underline{\mathfrak{F}}(\underline{D}) = \underline{\mathfrak{F}}_{b,\gamma}(V)$. The *admissible locus* is then defined as the subset over which $\underline{\mathfrak{F}}_{b,\gamma}(V)$ has slope zero. Then $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^{wa}$ and $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^a$ are open paracompact strictly \check{E} -analytic subspaces of $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}}^{\mathrm{an}}$ ([Hartl and Viehmann \[ibid.\]](#), Theorem 4.20).

The neutral admissible locus $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^{na} \subset \check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^a$ is defined by the condition that the images of $[b]$ and γ in $\pi_1(G)_{\Gamma}$ coincide.

We have the analog of the theorem of Colmez and Fontaine that “admissible implies weakly admissible”, see [Hartl \[2011\]](#), Theorem 2.5.3. In other words, we have an inclusion

$$\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^a \subseteq \check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^{wa}.$$

Furthermore, we have $\check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^a(L) = \check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^{wa}(L)$ for all finite field extensions L/\check{E} . If L is algebraically closed, weakly admissible does *not* imply admissible in general.

For an $\mathbb{F}[[\zeta]]$ -algebra B , complete and separated with respect to a bounded norm $|\cdot| : B \rightarrow [0, 1] \subset \mathbb{R}$ with $0 < |\zeta| < 1$, we consider the $\mathbb{F}((z))$ -algebra

$$B[[z, z^{-1}]] = \left\{ \sum_{i \in \mathbb{Z}} b_i z^i \mid b_i \in B, |b_i| |\zeta|^i \rightarrow 0 \ (i \rightarrow -\infty) \text{ for all } r > 0 \right\}$$

and the element $t_- = \prod_{i \in \mathbb{N}_0} \left(1 - \frac{\zeta^i}{z}\right) \in B[[z, z^{-1}]]$.

We want to define the *period morphism*

$$\check{\pi} : \check{\mathfrak{M}}^{\mathrm{an}} \rightarrow \check{\mathcal{H}}_{\mathbf{G}, \hat{Z}, b}^a$$

as in [Hartl and Viehmann \[2017\]](#), 6 where \mathfrak{M} is the Rapoport-Zink space associated with \hat{Z} and b . Let S be an affinoid, strictly \check{E} -analytic space and let $S \rightarrow \check{\mathfrak{M}}^{\mathrm{an}}$ be a morphism of

\check{E} -analytic spaces. With it we have to associate a morphism $S \rightarrow \check{\mathcal{H}}_{\mathbf{G}, \check{Z}}^{\text{an}}$. By construction of $\check{\mathfrak{M}}^{\text{an}}$ a morphism $S \rightarrow \check{\mathfrak{M}}^{\text{an}}$ is induced by a morphism from a quasi-compact admissible formal $\check{R}_{\check{Z}}$ -scheme \mathfrak{S} with $\mathfrak{S}^{\text{an}} = S$ to $\check{\mathfrak{M}}$. The latter corresponds to some $(\underline{\mathfrak{g}}, \bar{\delta}) \in \check{\mathfrak{M}}(\mathfrak{S})$. After an étale covering $\mathfrak{S}' = \text{Spf} B' \rightarrow \mathfrak{S}$ of admissible formal $\check{R}_{\check{Z}}$ -schemes there is a trivialization $\alpha : \underline{\mathfrak{g}}_{\mathfrak{S}'} \cong ((L^+ \mathbf{G})_{\mathfrak{S}'}, A\sigma^*)$ for some $A \in \mathbf{G}(B'[[z]][\frac{1}{z-\zeta}])$.

Denote by $\bar{\cdot}$ the reduction to $V(\zeta)$. Then the quasi-isogeny $\bar{\delta}$ induces via $\bar{\alpha}$ an element $\bar{\Delta} \in LG(\bar{B}')$. One then shows that there is a uniquely determined element $\Delta \in \mathbf{G}(B'[[z, z^{-1}]]_{[\frac{1}{z-\zeta}]})$ lifting $\bar{\Delta}$ with $\Delta A = b\sigma^*(\Delta)$ for A as above and b defining the Rapoport-Zink space. Consider the element

$$\gamma = \sigma^*(\Delta)A^{-1} \cdot \mathbf{G}(B'[\frac{1}{\xi}][[z - \zeta]]) \in \mathbf{G}(B'[\frac{1}{\xi}]((z - \zeta)))/\mathbf{G}(B'[\frac{1}{\xi}][[z - \zeta]]).$$

The boundedness of $\underline{\mathfrak{g}}$ then implies that γ factors through $\check{\mathcal{H}}_{\mathbf{G}, \check{Z}}^{\text{an}}$. It descends to a well-defined element of $\check{\mathcal{H}}_{\mathbf{G}, \check{Z}}^{\text{an}}(S)$ which is the image under the period morphism.

Proposition 7-2 (Hartl and Viehmann [2017], Prop. 6.9, 6.10). *The period morphism factors through the open \check{E} -analytic subspace $\check{\mathcal{H}}_{\mathbf{G}, \check{Z}, b}^a$ and induces an étale morphism*

$$\check{\pi} : \check{\mathfrak{M}}^{\text{an}} \rightarrow \check{\mathcal{H}}_{\mathbf{G}, \check{Z}, b}^a.$$

To describe the image of $\check{\pi}$ recall the tensor functors of (4-6) and (4-7). The construction of the σ -bundle $\check{\mathfrak{F}}_{b, \gamma}(V)$ of Definition 7-1 has a generalization to a σ -bundle $\check{\mathfrak{F}}_b(V)$ over $\check{\mathcal{H}}_{\mathbf{G}, \check{Z}}$ with fibers $\check{\mathfrak{F}}_{b, \gamma}(V)$. It further induces a canonical local system $\underline{\mathcal{V}}_b(V)$ of $\mathbb{F}_q((z))$ -vector spaces on $\check{\mathcal{H}}_{\mathbf{G}, \check{Z}, b}^a$ with $\underline{\mathcal{V}}_b(V)_{\bar{\gamma}} = \bar{\gamma}^* \check{\mathfrak{F}}_b(V)^{\tau}$ where we take invariants under the isomorphism τ defining the σ -bundle. Let $\omega_{b, \bar{\gamma}} : \text{Rep}_{\mathbb{F}_q((z))} G \rightarrow \text{FMod}_{\mathbb{F}_q((z))}$ be the fiber functor with $\omega_{b, \bar{\gamma}}(V) := \underline{\mathcal{V}}_b(V)_{\bar{\gamma}} = \bar{\gamma}^* \check{\mathfrak{F}}_b(V)^{\tau}$. Then, one of the main results of Hartl and Viehmann [ibid.] is the following description of the image of the period map. The condition that G be unramified can also be replaced by another, more technical, but possibly more general condition.

Theorem 7-3. *Assume that G is unramified.*

1. *The image $\check{\pi}(\check{\mathfrak{M}}^{\text{an}})$ of the period morphism is equal to the union of those connected components of $\check{\mathcal{H}}_{\mathbf{G}, \check{Z}, b}^a$ on which there is an $\mathbb{F}_q((z))$ -rational isomorphism $\beta : \omega^\circ \rightarrow \omega_{b, \bar{\gamma}}$.*
2. *The rational dual Tate module $\check{V}_{\underline{\mathfrak{g}}}$ of the universal local \mathbf{G} -shtuka $\underline{\mathfrak{g}}$ over $\check{\mathfrak{M}}^{\text{an}}$ descends to a tensor functor $\check{V}_{\underline{\mathfrak{g}}}$ from $\text{Rep}_{\mathbb{F}_q((z))} G$ to the category of local systems of $\mathbb{F}_q((z))$ -vector spaces on $\check{\pi}(\check{\mathfrak{M}}^{\text{an}})$. It carries a canonical $J_b(\mathbb{F}_q((z)))$ -linearization and is canonically $J_b(\mathbb{F}_q((z)))$ -equivariantly isomorphic to the tensor functor $\underline{\mathcal{V}}_b$.*

3. The tower of strictly \check{E} -analytic spaces $(\check{\mathfrak{M}}^K)_{K \subset \mathbf{G}(\mathbb{F}_q((z)))}$ is canonically isomorphic over $\check{\pi}(\check{\mathfrak{M}}^{\text{an}})$ in a Hecke and $J_b(\mathbb{F}_q((z)))$ -equivariant way to the tower of étale covering spaces of $\check{\pi}(\check{\mathfrak{M}}^{\text{an}})$ that is naturally associated with the tensor functor $\underline{\mathcal{U}}_b$.

References

- Greg W. Anderson (1993). “On Tate modules of formal t -modules”. *IMRN* 2, pp. 41–52. MR: [1203253](#) (cit. on p. [1445](#)).
- Esmail Arasteh Rad and Urs Hartl (2013). “Uniformizing the moduli stacks of global G -Shtukas”. To appear in *IMRN*. arXiv: [1302.6351](#) (cit. on pp. [1456](#), [1457](#)).
- (2014). “Local \mathbb{P} -shtukas and their relation to global \mathcal{G} -shtukas”. *Münster J. Math.* 7.2, pp. 623–670. MR: [3426233](#) (cit. on pp. [1445](#), [1447](#), [1450](#), [1451](#), [1453](#), [1455–1457](#)).
- François Bruhat and Jacques Tits (1972). “Groupes réductifs sur un corps local: I. Données radicielles valuées”. *Inst. Hautes Études Sci. Publ. Math.* 41, pp. 5–251. MR: [0327923](#) (cit. on p. [1444](#)).
- Alain Genestier (1996). “Espaces symétriques de Drinfeld”. *Astérisque* 234, pp. ii+124. MR: [1393015](#) (cit. on p. [1445](#)).
- Thomas Haines and Michael Rapoport (2008). “On parahoric subgroups”. appendix to [Pappas and Rapoport \[2008\]](#) (cit. on p. [1444](#)).
- Paul Hamacher and Eva Viehmann (Feb. 2017). “Irreducible components of minuscule affine Deligne-Lusztig varieties”. arXiv: [1702.08287](#) (cit. on pp. [1453](#), [1454](#)).
- Urs Hartl (2011). “Period spaces for Hodge structures in equal characteristic”. *Ann. of Math. (2)* 173.3, pp. 1241–1358. MR: [2800715](#) (cit. on p. [1461](#)).
- Urs Hartl and Wansu Kim (2016). “Local Shtukas, Hodge-Pink Structures and Galois Representations”. to appear in *Proceedings of the conference on “ t -motives: Hodge structures, transcendence and other motivic aspects”, BIRS, Banff, Canada 2009*, eds. G. Böckle, D. Goss, U. Hartl, M. Papanikolas, EMS (cit. on p. [1460](#)).
- Urs Hartl and Eva Viehmann (2011). “The Newton stratification on deformations of local \mathbf{G} -shtukas”. *J. Reine Angew. Math.* 656, pp. 87–129. MR: [2818857](#) (cit. on pp. [1445](#), [1448](#), [1449](#), [1453](#), [1455](#)).
- (2012). “Foliations in deformation spaces of local \mathbf{G} -shtukas”. *Adv. Math.* 229.1, pp. 54–78. MR: [2854170](#) (cit. on p. [1445](#)).
- (Dec. 2017). “The generic fiber of moduli spaces of bounded local G -shtukas”. arXiv: [1712.07936](#) (cit. on pp. [1447](#), [1449–1452](#), [1460–1462](#)).
- Xuhua He (2016). “Kottwitz-Rapoport conjecture on unions of affine Deligne-Lusztig varieties”. *Ann. Sci. Éc. Norm. Supér. (4)* 49.5, pp. 1125–1141. arXiv: [1408.5838](#). MR: [3581812](#) (cit. on p. [1453](#)).

- Xuhua He (2018). “Some results on affine Deligne–Lusztig varieties”. In: *Proc. of the International Congress of Mathematicians, Rio de Janeiro 2018*. Ed. by Boyan Sirakov, Paulo Ney de Souza, and Marcelo Viana. In this volume. World Scientific, pp. 1345–1366 (cit. on p. 1453).
- Robert E. Kottwitz (1985). “Isocrystals with additional structure”. *Compositio Math.* 56.2, pp. 201–220. MR: 809866 (cit. on p. 1445).
- (1997). “Isocrystals with additional structure. II”. *Compositio Math.* 109.3, pp. 255–339. MR: 1485921 (cit. on p. 1445).
- Vincent Lafforgue (Sept. 2012). “Chtoucas pour les groupes réductifs et paramétrisation de Langlands globale”. arXiv: 1209.5352 (cit. on p. 1454).
- Elena Mantovan (2004). “On certain unitary group Shimura varieties”. *Astérisque* 291. Variétés de Shimura, espaces de Rapoport–Zink et correspondances de Langlands locales, pp. 201–331. MR: 2074715 (cit. on p. 1457).
- Stephan Neupert (2016). “Foliations and the cohomology of moduli spaces of bounded global G -shtukas”. PhD thesis. TU Munich. arXiv: 1610.05935 (cit. on pp. 1445, 1450, 1455–1459).
- Sian Nie (2015). “Fundamental elements of an affine Weyl group”. *Math. Ann.* 362.1-2, pp. 485–499. MR: 3343886 (cit. on p. 1458).
- George Pappas and Michael Rapoport (2008). “Twisted loop groups and their affine flag varieties”. *Adv. Math.* 219.1. With an appendix by T. Haines and Rapoport, pp. 118–198. MR: 2435422 (cit. on pp. 1446, 1463).
- Michael Rapoport and Melanie Richartz (1996). “On the classification and specialization of F -isocrystals with additional structure”. *Compositio Math.* 103.2, pp. 153–181. MR: 1411570 (cit. on p. 1446).
- Michael Rapoport and Thomas Zink (1996). *Period spaces for p -divisible groups*. Vol. 141. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, pp. xxii+324. MR: 1393439 (cit. on pp. 1449, 1457).
- Timo Richarz (2016). “Affine Grassmannians and geometric Satake equivalences”. *IMRN* 2016, pp. 3717–3767. MR: 3544618 (cit. on p. 1446).
- Yakov Varshavsky (2004). “Moduli spaces of principal F -bundles”. *Selecta Math. (N.S.)* 10.1, pp. 131–166. MR: 2061225 (cit. on p. 1456).
- Eva Viehmann (2013). “Newton strata in the loop group of a reductive group”. *Amer. J. Math.* 135.2, pp. 499–518. MR: 3038719 (cit. on pp. 1453, 1454).
- (2014). “Truncations of level 1 of elements in the loop group of a reductive group”. *Ann. of Math. (2)* 179.3, pp. 1009–1040. MR: 3171757 (cit. on p. 1458).
- (Nov. 2015). “On the geometry of the Newton stratification”. arXiv: 1511.03156 (cit. on p. 1453).
- Eva Viehmann and Han Wu (June 2016). “Central leaves in loop groups”. To appear in *Math. Res. Letters*. arXiv: 1606.09444 (cit. on p. 1448).

Jean-Pierre Wintenberger (1997). “Propriétés du groupe tannakien des structures de Hodge p -adiques et toiseur entre cohomologies cristalline et étale”. *Ann. Inst. Fourier (Grenoble)* 47.5, pp. 1289–1334. MR: [1600395](#).

Received 2017-11-30.

EVA VIEHMANN
FAKULTÄT FÜR MATHEMATIK
TECHNISCHE UNIVERSITÄT MÜNCHEN – M11
BOLTZMANNSTR. 3
85748 GARCHING
GERMANY
viehmann@ma.tum.de

