

RANDOM BAND MATRICES

PAUL BOURGADE

Abstract

We survey recent mathematical results about the spectrum of random band matrices. We start by exposing the Erdős–Schlein–Yau dynamic approach, its application to Wigner matrices, and extension to other mean-field models. We then introduce random band matrices and the problem of their Anderson transition. We finally expose a method to obtain delocalization and universality in some sparse regimes, highlighting the role of quantum unique ergodicity.

Contents

1	Introduction	2777
2	Mean-field random matrices	2780
3	Random band matrices and the Anderson transition	2787
4	Quantum unique ergodicity and universality	2793

1 Introduction

This note explains the interplay between eigenvectors and eigenvalues statistics in random matrix theory, when the considered models are not of mean-field type, meaning that the interaction is short range and geometric constraints enter in the definition of the model.

If the range or strength of the interaction is small enough, it is expected that eigenvalues statistics will fall into the *Poisson universality class*, intimately related to the notion of independence. Another class emerged in the past fifty years for many correlated systems,

This work is supported by the NSF grant DMS#1513587.

MSC2010: 15B52.

Keywords: band matrices, delocalization, quantum unique ergodicity, Gaussian free field.

initially from calculations on random linear operators. This *random matrix universality class* was proposed by [Wigner \[1957\]](#), first as a model for stable energy levels of typical heavy nuclei. The models he introduced have since been understood to connect to integrable systems, growth models, analytic number theory and multivariate statistics (see e.g. [Deift \[2017\]](#)).

Ongoing efforts to understand universality classes are essentially of two types. First, *integrability* consists in finding possibly new statistics for a few, rigid models, with methods including combinatorics and representation theory. Second, *universality* means enlarging the range of models with random matrix statistics, through probabilistic methods. For example, the Gaussian random matrix ensembles are mean-field integrable models, from which local spectral statistics can be established for the more general Wigner matrices, by comparison, as explained in Section 1. For random operators with shorter range, no integrable models are known, presenting a major difficulty in understanding whether their spectral statistics will fall in the Poisson or random matrix class.

In Wigner's original theory, the eigenvectors play no role. However, their statistics are essential in view of a famous dichotomy of spectral behaviors, widely studied since Anderson's tight binding model [P. Anderson \[1958\]](#):

- (i) Poisson spectral statistics usually occur together with localized eigenstates,
- (ii) random matrix eigenvalue distributions should coincide with delocalization of eigenstates.

The existence of the localized phase has been established for the Anderson model in any dimension [Fröhlich and Spencer \[1983\]](#), but delocalization has remained elusive for all operators relevant in physics. An important question consists in proving extended states and GOE local statistics for one such model¹, giving theoretical evidence for conduction in solids. How localization implies Poisson statistics is well understood, at least for the Anderson model [Minami \[1996\]](#). In this note, we explain the proof of a strong notion of delocalization (quantum unique ergodicity), and how it implies random matrix spectral statistics, for the $1d$ random band matrix (RBM) model.

In this model, vertices are elements of $\Lambda = \llbracket 1, N \rrbracket^d$ ($d = 1, 2, 3$) and $H = (H_{ij})_{i,j \in \Lambda}$ have centered real entries, independent up to the symmetry $H_{ij} = H_{ji}$. The band width $W < N/2$ means

$$(1-1) \quad H_{ij} = 0 \text{ if } |i - j| > W,$$

where $|\cdot|$ is the periodic L^1 distance on Λ , and all non-trivial H_{ij} 's have a variance σ_{ij}^2 with the same order of magnitude, normalized by $\sum_j \sigma_{ij}^2 = 1$ for any $i \in \Lambda$. Mean-field

¹GOE eigenvalues statistics appear in Trotter's tridiagonal model [Trotter \[1984\]](#), which is clearly local, but the entries need varying variance adjusted to a specific profile.

models correspond to $W = N/2$. When $W \rightarrow \infty$, the empirical spectral measure of H converges to the semicircle distribution $d\rho_{sc}(x) = \frac{1}{2\pi}(4 - x^2)^{1/2}dx$.

It has been conjectured that the random band matrix model exhibits the localization-delocalization (and Poisson-GOE) transition at some critical band width $W_c(N)$ for eigenvalues in the bulk of the spectrum $|E| < 2 - \kappa$. The localized regime supposedly occurs for $W \ll W_c$ and delocalization for $W \gg W_c$, where

$$(1-2) \quad W_c = \begin{cases} N^{1/2} & \text{for } d = 1, \\ (\log N)^{1/2} & \text{for } d = 2, \\ O(1) & \text{for } d = 3. \end{cases}$$

This transition corresponds to localization length $\ell \approx W^2$ in dimension 1, $\ell \approx e^{W^2}$ in dimension 2.




Figure 1: Conjectural behavior of the RBM model for $d = 1$. For any eigenvalue $|\lambda_k| < 2 - \kappa$, the rescaled gap $N\rho_{sc}(\lambda_k)(\lambda_{k+1} - \lambda_k)$ converges to an exponential random variable for $W \ll N^{1/2}$, and the Gaudin GOE distribution for $W \gg N^{1/2}$. The associated eigenvector u_k is localized on $\ell \approx W^2$ sites for $W \ll N^{1/2}$, it is flat for $W \gg N^{1/2}$. Here $(Avf)(\alpha) = (2n)^{-1} \sum_{|i-\alpha|<n} f(i)$ where $1 \ll n \ll W^2$ is some averaging scale.

This review first explains universality techniques for mean-field models. We then state recent progress in proving the existence of the delocalized phase for the random band matrix model for $d = 1$, explaining how quantum unique ergodicity is proved by dynamics. We finally explain, at the heuristic level, a connection between quantum unique ergodicity for band matrices and the Gaussian free field, our main goal being to convince the reader that the transition exponents in (1-2) are natural.

For the sake of conciseness, we only consider the orthogonal symmetry class corresponding to random symmetric matrices with real entries. Analogous results hold in the complex Hermitian class.

2 Mean-field random matrices

2.1 Integrable model. The Gaussian orthogonal ensemble (GOE) consists in the probability density

$$(2-1) \quad \frac{1}{Z_N} e^{-\frac{N}{4} \text{Tr}(H^2)}$$

with respect to the Lebesgue measure on the set on $N \times N$ symmetric matrices. This corresponds to all entries being Gaussian and independent up to the symmetry condition, with off-diagonal entries $H_{ij} \sim N^{-1/2} \mathcal{N}(0, 1)$, and diagonal entries $H_{ii} \sim (N/2)^{-1/2} \mathcal{N}(0, 1)$.

Our normalization is chosen so that the eigenvalues $\lambda_1 \leq \dots \leq \lambda_N$ (with associated eigenvectors u_1, \dots, u_N) have a converging empirical measure: $\frac{1}{N} \sum_{k=1}^N \delta_{\lambda_k} \rightarrow d\rho_{\text{sc}}$ almost surely. A more detailed description of the spectrum holds at the microscopic scale, in the bulk and at the edge: there exists a translation invariant point process χ_1 [Mehta and Gaudin \[1960\]](#) and a distribution TW_1 (for [Tracy and Widom \[1994\]](#)) such that

$$(2-2) \quad \sum_{k=1}^N \delta_{N\rho_{\text{sc}}(E)(\lambda_k - E)} \rightarrow \chi_1,$$

$$(2-3) \quad N^{2/3}(\lambda_N - 2) \rightarrow \text{TW}_1,$$

in distribution. Note that χ_1 is independent of $E \in (-2 + \kappa, 2 - \kappa)$.

Concerning the eigenvectors, for any $\mathcal{O} \in O(N)$, from (2-1) the distributions of $\mathcal{O}^t H \mathcal{O}$ and H are the same, so that the eigenbasis $\mathbf{u} = (u_1, \dots, u_N)$ of H is Haar-distributed (modulo a sign choice) on $O(N)$: $\mathcal{O}\mathbf{u}$ has same distribution as \mathbf{u} . In particular, any u_k is uniform of the sphere $\mathcal{S}^{(N-1)}$, and has the same distribution as $\mathcal{N}/\|\mathcal{N}\|_2$ where \mathcal{N} is a centered Gaussian vector with covariance Id_N . This implies that for any deterministic sequences of indices $k_N \in \llbracket 1, N \rrbracket$ and unit vectors $\mathbf{q}_N \in \mathcal{S}^{(N-1)}$ (abbreviated k, \mathbf{q}), the limiting *Borel-Lévy law* holds:

$$(2-4) \quad N^{1/2} \langle u_{k_N}, \mathbf{q}_N \rangle \rightarrow \mathcal{N}(0, 1)$$

in distribution. This microscopic behavior can be extended to several projections being jointly Gaussian.

The fact that eigenvectors are extended can be formulated with less precision, and quantified in different manners. For example, for the GOE model, for any small $\varepsilon > 0$ and large $D > 0$, we have

$$(2-5) \quad \mathbb{P} \left(\|u_{k_N}\|_\infty \geq \frac{N^\varepsilon}{\sqrt{N}} \right) \leq N^{-D},$$

which we refer to as *delocalization* (the above N^ε can also be replaced by some logarithmic power).

Delocalization does not imply that the eigenvectors are flat in the sense of [Figure 1](#), as u_k could be supported on a small fraction of $\llbracket 1, N \rrbracket$. A strong notion of flat eigenstates was introduced by [Rudnick and Sarnak \[1994\]](#) for Riemannian manifolds: for any negatively curved and compact \mathfrak{M} with volume measure μ ,

$$(2-6) \quad \int_A |\psi_k(x)|^2 \mu(dx) \xrightarrow[k \rightarrow \infty]{} \int_A \mu(dx),$$

for any $A \subset \mathfrak{M}$. Here ψ_k is an eigenfunction (associated to the eigenvalue λ_k) of the Laplace-Beltrami operator, $0 \leq \lambda_1 \leq \dots \leq \lambda_k \leq \dots$ and $\|\psi_k\|_{L^2(\mu)} = 1$. This *quantum unique ergodicity* (QUE) notion strengthens the quantum ergodicity from [Shnirel'man \[1974\]](#), [Colin de Verdière \[1985\]](#), and [Zelditch \[1987\]](#), defined by an additional averaging on k and proved for a wide class of manifolds and deterministic regular graphs [Anantharaman and Le Masson \[2015\]](#) (see also [Brooks and Lindenstrauss \[2013\]](#)). QUE was rigorously proved for arithmetic surfaces, [Lindenstrauss \[2006\]](#), [Holowinsky \[2010\]](#), and [Holowinsky and Soundararajan \[2010\]](#). We will consider a probabilistic version of QUE at a local scale, for eigenvalues in the bulk of the spectrum ($\kappa \geq 0$ is small and fixed). By simple properties of the uniform measure on the unit sphere it is clear that the following version holds for the GOE: for any given (small) $\varepsilon > 0$ and (large) $D > 0$, for $N \geq N_0(\varepsilon, D)$, for any deterministic sequences $k_N \in \llbracket \kappa N, (1 - \kappa)N \rrbracket$ and $I_N \subset \llbracket 1, N \rrbracket$ (abbreviated k, I), $|I_N| > n$, we have

$$(2-7) \quad \mathbb{P} \left(\left| \sum_{\alpha \in I} (u_k(\alpha))^2 - \frac{1}{N} \right| > \frac{N^\varepsilon |I|^{1/2}}{N} \right) \leq N^{-D}.$$

We now consider the properties (2-2), (2-3) (2-4), (2-5), (3-3) for the following general model.

Definition 2.1 (Generalized Wigner matrices). *A sequence H_N (abbreviated H) of real symmetric centered random matrices is a generalized Wigner matrix if there exists $C, c > 0$ such that $\sigma_{ij}^2 := \text{Var}(H_{ij})$ satisfies*

$$(2-8) \quad c \leq N\sigma_{ij}^2 \leq C \text{ for all } N, i, j \text{ and } \sum_j \sigma_{ij}^2 = 1 \text{ for all } i.$$

We also assume subgaussian decay of the distribution of $\sqrt{N} H_{ij}$, uniformly in i, j, N , for convenience (this could be replaced by a finite high moment assumption).

2.2 Eigenvalues universality. The second constraint in (2-8) imposes the macroscopic behavior of the limiting spectral measure: $\frac{1}{N} \sum_{k=1}^N \delta_{\lambda_i} \rightarrow d\rho_{\text{sc}}$ for all generalized Wigner matrices. This convergence to the semicircle distribution was strengthened up to optimal polynomial scale, thanks to an advanced diagrammatic analysis of the resolvent of H .

Theorem 2.2 (Rigidity of the spectrum Erdős, Yau, and Yin [2012b]). *Let H be a generalized Wigner matrix as in Definition 2.1. Define $\hat{k} = \min(k, N+1-k)$ and γ_k implicitly by $\int_{-2}^{\gamma_k} d\rho_{\text{sc}} = \frac{k}{N}$. Then for any $\varepsilon > 0$, $D > 0$ there exists N_0 such that for $N > N_0$ we have*

$$(2-9) \quad \mathbb{P} \left(|\lambda_k - \gamma_k| > N^{-\frac{2}{3} + \varepsilon} (\hat{k})^{-\frac{1}{3}} \right) \leq N^{-D}.$$

Given the above scale of fluctuations, a natural problem consists in the limiting distribution. In particular, the (Wigner-Dyson-Mehta) conjecture states that (2-2) holds for random matrices way beyond the integrable GOE class. It has been proved in a series of works in the past years, with important new techniques based on the Harish-Chandra-Itzykson-Zuber integral Johansson [2001] (in the special case of Hermitian symmetry class), the dynamic interpolation through Dyson Brownian motion Erdős, Schlein, and Yau [2011] and the Lindeberg exchange principle Tao and Vu [2011]. The initial universality statements for general classes required an averaging over the energy level E Erdős, Schlein, and Yau [2011] or the first four moments of the matrix entries to match the Gaussian ones Tao and Vu [2011].

We aim at explaining the dynamic method which was applied in a remarkable variety of settings. For example, GOE local eigenvalues statistics holds for generalized Wigner matrices.

Theorem 2.3 (Fixed energy universality Bourgade, Erdős, Yau, and Yin [2016]). *The convergence (2-2) holds for generalized Wigner matrices.*

The key idea for the proof, from Erdős, Schlein, and Yau [2011], is interpolation through matrix Dyson Brownian motion (or its Ornstein Uhlenbeck version)

$$(2-10) \quad dH_t = \frac{1}{\sqrt{N}} dB_t - \frac{1}{2} H_t dt$$

with initial condition $H_0 = H$, where $(B_{ij})_{i < j}$ and $(B_{ii}/\sqrt{2})_i$ are independent standard Brownian motions. The GOE measure (2-1) is the equilibrium for these dynamics. The proof proceeds in two steps, in which the dynamics $(H_t)_{t \geq 0}$ is analyzed through complementary viewpoints. One relies on the repulsive eigenvalues dynamics, the other on the matrix structure. Both steps require some a priori knowledge on eigenvalues density, such as Theorem 2.2.

First step: relaxation. For any $t \geq N^{-1+\varepsilon}$, (2-2) holds: $\sum_{k=1}^N \delta_{N\rho_{\text{sc}}(E)(\lambda_k(t)-E)} \rightarrow \chi_1$, where we denote $\lambda_1(t) \leq \dots \leq \lambda_k(t)$ the eigenvalues of H_t . The proof relies on the Dyson Brownian motion for the eigenvalues dynamics [Dyson \[1962\]](#), given by

$$(2-11) \quad d\lambda_k(t) = \frac{d\widetilde{B}_k(t)}{\sqrt{N}} + \left(\frac{1}{N} \sum_{\ell \neq k} \frac{1}{\lambda_k(t) - \lambda_\ell(t)} - \frac{1}{2} \lambda_k(t) \right) dt$$

where the $\widetilde{B}_k/\sqrt{2}$'s are standard Brownian motions. Consider the dynamics with a different initial condition $x_1(0) \leq \dots \leq x_N(0)$ given by the eigenvalues of a GOE matrix, and by taking the difference between these two equations we observe that $\delta_\ell(t) := e^{t/2}(x_\ell(t) - \lambda_\ell(t))$ satisfy an integral equation of parabolic type [Bourgade, Erdős, Yau, and Yin \[2016\]](#), namely

$$(2-12) \quad \partial_t \delta_\ell(t) = \sum_{k \neq \ell} b_{k\ell}(t)(\delta_k(t) - \delta_\ell(t)), \quad b_{k\ell}(t) = \frac{1}{N(x_\ell(t) - x_k(t))(\lambda_\ell(t) - \lambda_k(t))}.$$

From [Theorem 2.2](#), in the bulk of the spectrum we expect that $b_{k\ell}(t) \approx N/(k - \ell)^2$, so that Hölder regularity holds for $t \gg N^{-1}$: $\delta_k(t) = \delta_{k+1}(t)(1 + o(1))$, meaning $\lambda_{k+1}(t) - \lambda_k(t) = y_{k+1}(t) - y_k(t) + o(N^{-1})$. Gaps between the λ_k 's and x_k 's therefore become identical, hence equal to the GOE gaps as the law of $y_{k+1}(t) - y_k(t)$ is invariant in time. In fact, an equation similar to (2-12) previously appeared in the first proof of GOE gap statistics for generalized Wigner matrices [Erdős and Yau \[2015\]](#), emerging from a Helffer-Sjöstrand representation instead of a probabilistic coupling. [Theorem 2.3](#) requires a much more precise analysis of (2-12) [Bourgade, Erdős, Yau, and Yin \[2016\]](#) and [Landon, Sosoe, and Yau \[2016\]](#), but the conceptual picture is clear from the above probabilistic coupling of eigenvalues.

Relaxation after a short time can also be understood by functional inequalities for relative entropy [Erdős, Schlein, and Yau \[2011\]](#) and [Erdős, Yau, and Yin \[2012a\]](#), a robust method which also gives GOE statistics when averaging over the energy level E . In the special case of the Hermitian symmetry class, relaxation also follows from explicit formulas for the eigenvalues density at time t [Johansson \[2001\]](#), [Erdős, Péché, Ramírez, Schlein, and Yau \[2010\]](#), and [Tao and Vu \[2011\]](#).

Second step: density. For any $t \leq N^{-\frac{1}{2}-\varepsilon}$,

$$\sum_{k=1}^N \delta_{N\rho_{\text{sc}}(E)(\lambda_k(t)-E)} \quad \text{and} \quad \sum_{k=1}^N \delta_{N\rho_{\text{sc}}(E)(\lambda_k(0)-E)}$$

have the same distribution at leading order. This step can be proved by a simple Itô lemma based on the matrix evolution [Bourgade and Yau \[2017\]](#), which takes a particularly simple

form for Wigner matrices (i.e. $\sigma_{ij}^2 = N^{-1} + N^{-1}\mathbb{1}_{i=j}$). It essentially states that for any smooth function $F(H)$ we have

$$(2-13) \quad \mathbb{E} F(H_t) - \mathbb{E} F(H_0) = O(tN^{1/2})$$

$$\sup_{i \leq j, 0 \leq s \leq t} \mathbb{E} \left((N^{3/2} |H_{ij}(s)|^3 + \sqrt{N} |H_{ij}(s)|) |\partial_{ij}^3 F(H_s)| \right)$$

where $\partial_{ij} = \partial_{H_{ij}}$. In particular, if F is stable in the sense that $\partial_{ij}^3 F = O(N^\varepsilon)$ with high probability (this is known for functions encoding the microscopic behavior thanks to the a-priori rigidity estimates from [Theorem 2.2](#)), then invariance of local statistics holds up to time $N^{-\frac{1}{2}-\varepsilon}$.

Invariance of local spectral statistics in matrix neighborhoods has also been proved by other methods, for example by a reverse heat flow when the entries have a smooth enough density [Erdős, Schlein, and Yau \[2011\]](#), or the Lindeberg exchange principle [Tao and Vu \[2011\]](#) for matrices with moments of the entries coinciding up to fourth moment.

2.3 Eigenvectors universality. Eigenvalues rigidity (2-9) was an important estimate for the proof of [Theorem 2.3](#). Similarly, to understand the eigenvectors distribution, one needs to first identify their natural fluctuation scale. By analysis of the resolvent of H , the following was first proved when \mathbf{q} is an element from the canonical basis [Erdős, Schlein, and Yau \[2009\]](#) and [Erdős, Yau, and Yin \[2012b\]](#), and extended to any direction.

Theorem 2.4 (Isotropic delocalization [Knowles and Yin \[2013b\]](#) and [Bloemendal, Erdős, Knowles, Yau, and Yin \[2014\]](#)). *For any sequence of generalized Wigner matrices, $\varepsilon, D > 0$, there exists $N_0(\varepsilon, D)$ such that for any $N \geq N_0$ deterministic k and unit vector \mathbf{q} ,*

$$\mathbb{P} \left(\langle u_k, \mathbf{q} \rangle \geq N^{-\frac{1}{2}+\varepsilon} \right) \leq N^{-D}.$$

The more precise fluctuations (2-4) were proved by the Lindeberg exchange principle in [Knowles and Yin \[2013a\]](#) and [Tao and Vu \[2012\]](#), under the assumption of the first four (resp. two) moments of H matching the Gaussian ones for eigenvectors associated to the spectral bulk (resp. edge). This Lévy-Borel law holds without these moment matching assumptions, and some form of quantum unique ergodicity comes with it.

Theorem 2.5 (Eigenvectors universality and weak QUE [Bourgade and Yau \[2017\]](#)). *For any sequence of generalized Wigner matrices, and any deterministic k and unit vector \mathbf{q} , the convergence (2-4) is true.*

Moreover, for any $\varepsilon > 0$ there exists $D > 0$ such that (3-3) holds.

The above statement is a *weak* form of QUE, holding for some small $D = D(\varepsilon)$ although it should be true for any large $D > 0$. Section 3 will show a strong form of QUE for some band matrices.

The proof of [Theorem 2.5](#) follows the dynamic idea already described for eigenvalues, by considering the evolution of eigenvectors through (2-10). The *density* step is similar: with (2-13) one can show that the distribution of $\sqrt{N}\langle u_k(t), \mathbf{q} \rangle$ is invariant up to time $t \leq N^{-\frac{1}{2}-\varepsilon}$. The *relaxation step* is significantly different from the coupling argument described previously. The eigenvectors dynamics are given by


$$du_k = \frac{1}{\sqrt{N}} \sum_{\ell \neq k} \frac{d\tilde{B}_{k\ell}}{\lambda_k - \lambda_\ell} u_\ell - \frac{1}{2N} \sum_{\ell \neq k} \frac{dt}{(\lambda_k - \lambda_\ell)^2} u_k,$$

where the $\tilde{B}_{k\ell}$'s are independent standard Brownian motions, and most importantly independent from the \tilde{B}_k 's from (2-11). This eigenvector flow was computed in the context of Brownian motion on ellipsoids [Norris, Rogers, and Williams \[1986\]](#), real Wishart processes [Bru \[1989\]](#), and for GOE/GUE in [G. W. Anderson, Guionnet, and Zeitouni \[2010\]](#).

Due to its complicated structure and high dimension, this eigenvector flow had not been analyzed. Surprisingly, these dynamics can be reduced to a multi-particle random walk in a dynamic random environment. More precisely, a configuration η consists in d points of $\llbracket 1, N \rrbracket$, with possible repetition. The number of particles at site x is η_x . A configuration obtained by moving a particle from i to j is denoted η^{ij} . The main observation from [Bourgade and Yau \[2017\]](#) is as follows. First denote $z_k = \sqrt{N}\langle \mathbf{q}, u_k \rangle$, which is random and time dependent. Then associate to a configuration η with j_k points at i_k , the renormalized moments observables (the \mathcal{N}_{i_k} are independent Gaussians) conditionally to the eigenvalues path,

$$(2-14) \quad f_{t,\lambda}(\eta) = \mathbb{E} \left(\prod_k z_{i_k}^{2j_k} \mid \lambda \right) / \mathbb{E} \left(\prod_k \mathcal{N}_{i_k}^{2j_k} \right).$$

Then $f_{t,\lambda}$ satisfies the parabolic partial differential equation



$$(2-15) \quad \partial_t f_{t,\lambda}(\eta) = \mathcal{B}(t) f_{t,\lambda}(\eta)$$

where

$$\mathcal{B}(t) f(\eta) = \frac{1}{N} \sum_{i \neq j} 2\eta_i (1 + 2\eta_j) \frac{f(\eta^{ij}) - f(\eta)}{(\lambda_i(t) - \lambda_j(t))^2}.$$

As shown in the above drawing, the generator $\mathcal{B}(t)$ corresponds to a random walk on the space of configurations η , with time-dependent rates given by the eigenvalues dynamics. This equation is parabolic and by the scale argument explained for (2-12), $f_{t,\lambda}$ becomes locally constant (in fact, equal to 1 by normalization constraint) for $t \geq N^{-1+\varepsilon}$. This Hölder regularity is proved by a maximum principle.

2.4 Other models. The described dynamic approach applies beyond generalized Wigner matrices. We do not attempt to give a complete list of applications of this method. Below are a few results.

- (i) Wigner-type matrices designate variations of Wigner matrices with non centered H_{ii} 's [Lee, Schnell, Stetler, and Yau \[2015\]](#), or the normalization constraint in (2-8) not satisfied (the limiting spectral measure may differ from semicircular) [Ajanki, Erdős, and Kruger \[2017\]](#), or the H_{ij} 's non-centered and correlated [Ajanki, Erdős, and Kruger \[2018\]](#) and [Erdős, Kruger, and Schroder \[2017\]](#). In all cases, GOE bulk statistics is known.
- (ii) For small mean-field perturbations of diagonal matrices (the Rosenzweig-Porter model), GOE statistics [Landon, Sosoe, and Yau \[2016\]](#) occur with localization [Benigni \[2017\]](#) and [von Soosten and Warzel \[2017\]](#). We refer to [Facoetti, Vivo, and Biroli \[2016\]](#) for the physical meaning of this unusual regime.
- (iii) Random graphs also have bulk or edge GOE statistics when the connectivity grows fast enough with N , as proved for example for the Erdős–Renyi [Erdős, Knowles, Yau, and Yin \[2012\]](#), [Landon, Huang, and Yau \[2015\]](#), [Huang, Landon, and Yau \[2017\]](#), and [Lee and Schnell \[2015\]](#) and uniform d -regular models [Bauerschmidt, Huang, Knowles, and Yau \[2017\]](#). Eigenvectors statistics are also known to coincide with the GOE for such graphs [Bourgade, Huang, and Yau \[2017\]](#).
- (iv) The convolution model $D_1 + U^*D_2U$, where D_1, D_2 are diagonal and U is uniform on $O(N)$, appears in free probability theory. Its empirical spectral measure is understood up to the optimal scale [Bao, Erdős, and Schnell \[2017\]](#), and GOE bulk statistics were proved in [Che and Landon \[2017\]](#).
- (v) For β -ensembles, the external potential does not impact local statistics, a fact first shown when $\beta = 1, 2, 4$ (the classical invariant ensembles) by asymptotics of orthogonal polynomials [Bleher and Its \[1999\]](#), [Deift \[1999\]](#), [Deift and Goev \[2009\]](#), [Lubinsky \[2009\]](#), and [L. Pastur and M. Shcherbina \[1997\]](#). The dynamics approach extended this result to any β [Bourgade, Erdős, and Yau \[2014a,b\]](#). Other methods

based on sparse models [Krishnapur, Rider, and Virág \[2016\]](#) and transport maps [Bekerman, Figalli, and Guionnet \[2015\]](#) and [M. Shcherbina \[2014\]](#) were also applied to either β -ensembles or multimatrix models [Figalli and Guionnet \[2016\]](#).

- (vi) Extremal statistics. The smallest gaps in the spectrum of Gaussian ensembles and Wigner matrices have the same law [Bourgade \[2018\]](#), when the matrix entries are smooth. The relaxation step (2-12) was quantified with an optimal error so that the smallest spacing scale ($N^{-4/3}$ in the GUE case [Arous and Bourgade \[2011\]](#)) can be perceived.

The above models are mean-field, a constraint inherent to the dynamic proof strategy. Indeed, the density step requires the matrix entries to fluctuate substantially: lemmas of type (2-13) need a constant variance of the entries along the dynamics (2-10).

3 Random band matrices and the Anderson transition

In the Wigner random matrix model, the entries, which represent the quantum transition rates between two quantum states, are all of comparable size. More realistic models involve geometric structure, as typical quantum transitions only occur between nearby states. In this section we briefly review key results for Anderson and band matrix models.

3.1 Brief and partial history of the random Schrödinger operator. Anderson's random Schrödinger operator [P. Anderson \[1958\]](#) on \mathbb{Z}^d describes a system with spatial structure. It is of type

$$(3-1) \quad H_{\text{RS}} = \Delta + \lambda V$$

where Δ is the discrete Laplacian and the random variables $V(x)$, $x \in \mathbb{Z}^d$, are i.i.d and centered with variance 1. The parameter $\lambda > 0$ measures the strength of the disorder. The spectrum of H_{RS} is supported on $[-2d, 2d] + \lambda \text{supp}(\mu)$ where μ is the distribution of $V(0)$

Amongst the many mathematical contributions to the analysis of this model, Anderson's initial motivation (localization, hence the suppression of electron transport due to disorder) was proved rigorously by [Fröhlich and Spencer \[1983\]](#) by a multiscale analysis: localization holds for strong disorder or at energies where the density of states $\rho(E)$ is very small (localization for a related one-dimensional model was previously proved by [Gold-sheid, S. A. Molchanov, and L. Pastur \[1977\]](#)). An alternative derivation was given in [Aizenman and S. Molchanov \[1993\]](#), who introduced a fractional moment method. From the scaling theory of localization [Abraham, P. W. Anderson, Licciardello, and Ramakrishnan \[1979\]](#), extended states supposedly occur in dimensions $d \geq 3$ for λ small enough, while eigenstates are only marginally localized for $d = 2$.

Unfortunately, there has been no progress in establishing the delocalized regime for the random Schrödinger operator on \mathbb{Z}^d . The existence of absolutely continuous spectrum (related to extended states) in the presence of substantial disorder is only known when \mathbb{Z}^d is replaced by homogeneous trees [Klein \[1994\]](#).

These results and conjecture were initially for the Anderson model in infinite volume. If we denote H_{RS}^N the operator (3-1) restricted to the box $\llbracket -N/2, N/2 \rrbracket^d$ with periodic boundary conditions, its spectrum still lies on a compact set and one expects that the bulk eigenvalues in the microscopic scaling (i.e. multiplied by N^d) converge to either Poisson or GOE statistics (H_{RS}^N corresponds to GOE rather than GUE because it is a real symmetric matrix). Minami proved Poisson spectral statistics from exponential decay of the resolvent [Minami \[1996\]](#), in cases where localization in infinite volume is known. For H_{RS}^N , not only is the existence of delocalized states in dimension three open, but also there is no clear understanding about how extended states imply GOE spectral statistics.

3.2 Random band matrices: analogies, conjectures, heuristics. The band matrix model we will consider was essentially already defined around (1-1). In addition, in the following we will assume subgaussian decay of the distribution of $W^{\frac{d}{2}} H_{ij}$, uniformly in i, j, N , for convenience (this could be replaced by a finite high moment assumption).

Although random band matrices and the random Schrödinger operator (3-1) are different, they are both local (their matrix elements H_{ij} vanish when $|i - j|$ is large). The models are expected to have the same properties when

$$(3-2) \quad \lambda \approx \frac{1}{W}.$$

For example, eigenvectors for the Anderson model in one dimension are proved to decay exponentially fast with a localization length proportional to λ^2 , in agreement with the analogy (3-2) and the [Equation \(1-2\)](#) when $d = 1$. For $d = 2$, it is conjectured that all states are localized with a localization length of order $\exp(W^2)$ for band matrices, $\exp(\lambda^{-2})$ for the Anderson model, again coherently with (3-2) and (1-2). For some mathematical justification of the analogy (3-2) from the point of view of perturbation theory, we refer to [Spencer \[2012, Appendix 4.11\]](#).

The origins of [Equation \(1-2\)](#) first lie on numerical evidence, at least for $d = 1$. In [Casati, Molinari, and Izrailev \[1990Apr\]](#) it was observed, based on computer simulations, that the bulk eigenvalue statistics and eigenvector localization length of $1d$ random band matrices are essentially a function of W^2/N , with the sharp transition as described before (1-2). Fyodorov and Mirlin gave the first theoretical explanation for this transition [Fyodorov and Mirlin \[1991\]](#). They considered a slightly different ensemble with complex Gaussian entries decaying exponentially fast at distance greater than W from the diagonal.

Based on a non-rigorous supersymmetric approach [Efetov \[1997\]](#), they approximate relevant random matrix statistics with expectations for a σ -model approximation, from which a saddle point method gives the localization/delocalization transition for $W \approx \sqrt{N}$. Their work also gives an estimate on the localization length ℓ , anywhere in the spectrum [Fyodorov and Mirlin \[1991\]](#), equation (19): it is expected that at energy level E (remember our normalization $\sum_j \sigma_{ij}^2 = 1$ for H so that the equilibrium measure is ρ_{sc}),

$$\ell \approx W^2(4 - E^2).$$

Finally, heuristics for localization/delocalization transition exponents follow from the conductance fluctuations theory developed by [Thouless \[1977\]](#), based on scaling arguments. For a discussion of mathematical aspects of the Thouless criterion, see [Spencer \[2012, 2010\]](#), and [Wang \[1992, Section III\]](#) for some rigorous scaling theory of localization. This criterion was introduced in the context of Anderson localisation, and was applied in [Sodin \[2010, 2014\]](#) to $1d$ band matrices, including at the edge of the spectrum, in agreement with the prediction from [Fyodorov and Mirlin \[1991\]](#). A different heuristic argument for (1-2) is given in Section 3, for any dimension in the bulk of the spectrum.

3.3 Results. The density of states ($\mathbb{E}(N^{-1} \sum_k \delta_{\lambda_k})$) of properly scaled random band matrices in dimension 1 converges to the semicircular distribution for any $W \rightarrow \infty$, as proved in [Bogachev, S. A. Molchanov, and L. A. Pastur \[1991\]](#). This convergence was then strengthened and fluctuations around the semicircular law were studied in [Guionnet \[2002\]](#), [G. W. Anderson and Zeitouni \[2006\]](#), [Jana, Saha, and Soshnikov \[2016\]](#), and [Li and Soshnikov \[2013\]](#) by the method of moments, at the macroscopic scale.

Interesting transitions extending the microscopic one (1-2) are supposed to occur at mesoscopic scales η , giving a full phase diagram in (η, W) . The work [Erdős and Knowles \[2015\]](#) rigorously analyzed parts of this diagram by studying linear statistics in some mesoscopic range and in any dimension, also by a moment-based approach.

The microscopic scale transitions (1-2) are harder to understand, but recent progress allowed to prove the existence of localization and delocalization for some polynomial scales in W . These results are essentially of four different types: (i) the localization side for general models, (ii) localization and delocalization for specific Gaussian models, (iii) delocalization for general models. Finally, (iv) the edge statistics are fully understood by the method of moments. Unless otherwise stated, all results below are restricted to $d = 1$.

(i) *Localization for general models.* A seminal result in the analysis of random band matrices is the following estimate on the localization scale. For simplicity one can assume that the entries of H are i.i.d. Gaussian, but the method from [Schenker \[2009\]](#) allows to treat more general distributions.

Theorem 3.1 (The localization regime for band matrices [Schenker \[2009\]](#)). *Let $\mu > 8$. There exists $\tau > 0$ such that for large enough N , for any $\alpha, \beta \in \llbracket 1, N \rrbracket$ one has*

$$\mathbb{E} \left(\sup_{1 \leq k \leq N} |u_k(\alpha)u_k(\beta)| \right) \leq W^\tau e^{-\frac{|\alpha-\beta|}{W^\mu}}.$$

Localization therefore holds simultaneously for all eigenvectors when $W \ll N^{1/8}$, which was improved to $W \ll N^{1/7}$ in [Peled, Schenker, Shamis, and Sodin \[2017\]](#) for some specific Gaussian model described below.

(ii) *Gaussian models with specific variance profile and supersymmetry.* For some Gaussian band matrices, the supersymmetry (SUSY) technique gives a purely analytic technique towards spectral properties. This approach has first been developed by physicists [Efetov \[1997\]](#). A rigorous supersymmetry method started with the expected density of states on arbitrarily short scales for a $3d$ band matrix ensemble [Disertori, Pinson, and Spencer \[2002\]](#), extended to $2d$ in [Disertori and Lager \[2017\]](#) (see [Spencer \[2012\]](#) for much more about the mathematical aspects of SUSY). More recently, the work [T. Shcherbina \[2014b\]](#) proved local GUE local statistics for $W \geq cN$, and delocalization was obtained in a strong sense for individual eigenvectors, when $W \gg N^{6/7}$ and the first four moments of the matrix entries match the Gaussian ones [Bao and Erdős \[2017\]](#). These recent rigorous results assume complex entries and hold for $|E| < \sqrt{2}$, for a block-band structure of the matrix with a specific variance profile.

We briefly illustrate the SUSY method for moments of the characteristic polynomial: remarkably, this is currently the only observable for which the transition at $W \approx \sqrt{N}$ was proved. Consider a matrix H whose entries are complex centered Gaussian variables such that

$$\mathbb{E}(H_{ij}H_{\ell k}) = \mathbb{1}_{i=k, j=\ell} J_{ij} \text{ where } J_{ij} = (-W^2\Delta + 1)_{ij}^{-1},$$

and Δ is the discrete Laplacian on $\llbracket 1, N \rrbracket$ with periodic boundary condition. The variance J_{ij} is exponentially small for $|i - j| > W^{1+\varepsilon}$, so that H can be considered a random band matrix with band width W . Define

$$F_2(E_1, E_2) = \mathbb{E} (\det(E_1 - H) \det(E_2 - H)), \quad D_2 = F_2(E, E).$$

Theorem 3.2 (Transition at the level of characteristic polynomials [T. Shcherbina \[2014a\]](#) and [M. Shcherbina and T. Shcherbina \[2017\]](#)). *For any $E \in (-2, 2)$ and $\varepsilon > 0$, we have*

$$\lim_{N \rightarrow \infty} (D_2)^{-1} F_2 \left(E + \frac{x}{N\rho_{\text{sc}}(E)}, E - \frac{x}{N\rho_{\text{sc}}(E)} \right) = \begin{cases} \frac{\sin(2\pi x)}{2\pi x} & \text{if } N^\varepsilon < W < N^{\frac{1}{2}-\varepsilon} \\ 1 & \text{if } N^{\frac{1}{2}+\varepsilon} < W < N \end{cases}.$$

Unfortunately, currently the local eigenvalues statistics cannot be identified from moments of characteristic polynomials: they require ratios which are more difficult to analyze by the SUSY method.

We briefly mention the key steps of the proof of [Theorem 3.2](#). First, an integral representation for F_2 is obtained by integration over Grassmann variables. These variables give convenient formulas for the product of characteristic polynomials: they allow to express the determinant as Gaussian-type integral. Integrate over the Grassmann variables then gives an integral representation (in complex variables) of the moments of interest. More precisely, the Gaussian representation for $F_2 \left(E + \frac{x}{N\rho_{sc}(E)}, E - \frac{x}{N\rho_{sc}(E)} \right)$, from [T. Shcherbina \[2014a\]](#), is

$$\frac{1}{(2\pi)^N} \frac{1}{\det J^2} \int e^{-\frac{W^2}{2} \sum_{j=-n+1}^n \text{Tr}(X_j - X_{j-1})^2 - \frac{1}{2} \sum_{j=-n}^n \text{Tr}(X_j + \frac{i\Delta_E}{2} + i \frac{\Delta_x}{N\rho_{sc}(E)})^2} \prod_{j=-n}^n \det(X_j - i\Delta_E/2) dX_j,$$

where $N = 2n + 1$, $\Delta_E = \text{diag}(E, E)$, $\Delta_x = \text{diag}(x, -x)$, and dX_j is the Lebesgue measure on 2×2 Hermitian matrices. This form of the correlation of characteristic polynomial is then analyzed by steepest descent. Analogues of the above representation hold in any dimension, where the matrices X_j, X_k , are coupled in a quadratic way when k and j are neighbors in \mathbb{Z}^d , similarly to the Gaussian free field.

Finally, based on their integral representations, it is expected that random band matrices behave like σ -models, which are used by physicists to understand complicated statistical mechanics systems. We refer to the recent work [M. Shcherbina and T. Shcherbina \[2018\]](#) for rigorous results in this direction.

(iii) Delocalization for general models. Back to general models with no specific distribution of the entries (except sufficient decay of the distribution, for example subgaussian), the first delocalization results for random band matrices relied on a difficult analysis of their resolvent.

For example, the Green’s function was controlled down to the scale W^{-1} in [Erdős, Yau, and Yin \[2012a\]](#), implying that the localization length of all eigenvectors is at least W . Analysis of the resolvent also gives full delocalization for most eigenvectors, for W large enough. In the theorem below, We say that an eigenvector u_k is subexponentially localized at scale ℓ if there exists $\varepsilon > 0$, $I \subset \llbracket 1, N \rrbracket$, $|I| \leq \ell$, such that $\sum_{\alpha \notin I} |u_k(\alpha)|^2 < e^{-N^\varepsilon}$.

Theorem 3.3 (Delocalized regime on average [Erdős, Knowles, Yau, and Yin \[2013\]](#)). *Assume $W \gg N^{4/5}$ and $\ell \ll N$. Then the fraction of eigenvectors subexponentially localized on scale ℓ vanishes as $N \rightarrow \infty$, with large probability.*

This result when $W \geq N^{6/7}$ was previously obtained in [Erdős and Knowles \[2011\]](#), and similar statements were proved in higher dimension.

Delocalization was recently proved without averaging, together with eigenvalues statistics and flatness of individual eigenvectors. The main new ingredient is that quantum unique ergodicity is a convenient delocalization notion, proved by dynamics.

To simplify the statement below, assume that H is a Gaussian-divisible, in the sense that $\sqrt{W}H_{ij}$ is the sum of two independent random variables, $X + \mathcal{N}(0, c)$, where c is an arbitrary small constant (the result holds for more general entries).

Theorem 3.4 (Delocalized regime [Bourgade, Yau, and Yin \[2018\]](#)). *Assume $W \gg N^{3/4+a}$ for some $a > 0$. Let $\kappa > 0$ be fixed.*

- (a) *For any $E \in (-2 + \kappa, 2 - \kappa)$ the eigenvalues statistics at energy level E converge to the GOE, as in (2-2).*
- (b) *The bulk eigenvectors are delocalized: for any (small) $\varepsilon > 0$, (large) $D > 0$, for $N \geq N_0(\varepsilon, D, \kappa)$ and $k \in \llbracket \kappa N, (1 - \kappa)N \rrbracket$, we have*

$$\mathbb{P} \left(\|u_k\|_\infty > N^{-\frac{1}{2} + \varepsilon} \right) < N^{-D}.$$

- (c) *The bulk eigenvectors are flat on any scale greater than W . More precisely, for any given (small) $\varepsilon > 0$ and (large) $D > 0$, for $N \geq N_0(\varepsilon, D)$, for any deterministic $k \in \llbracket \kappa N, (1 - \kappa)N \rrbracket$ and interval $I \subset \llbracket 1, N \rrbracket$, $|I_N| > W$, we have*

$$(3-3) \quad \mathbb{P} \left(\left| \sum_{\alpha \in I} (u_k(\alpha))^2 - \frac{1}{N} \right| > N^{-a + \varepsilon} \frac{|I|}{N} \right) \leq N^{-D}.$$

A strong form of QUE similar to (c) holds for random d -regular graphs [Bauerschmidt, Huang, and Yau \[2016\]](#), the proof relying on exchangeability. For models with geometric constraints, other ideas are explained in the next section.

[Theorem 3.4](#) relies on a mean-field reduction strategy initiated in [Bourgade, Erdős, Yau, and Yin \[2017\]](#), and an extension of the dynamics (2-15) to observables much more general than (2-14), as explained in the next section. New ingredients compared to 3.4 are (a) quantum unique ergodicity for mean-field models after Gaussian perturbation, in a strong sense, (b) estimates on the resolvent of the band matrix at the (almost macroscopic) scale $N^{-\varepsilon}$.

The current main limitation of the method to approach the transition $W_c = N^{1/2}$ comes from (b). These resolvent estimates are obtained by intricate diagrammatics developed in a series of previous works including [Erdős, Knowles, Yau, and Yin \[2013\]](#), and currently only proved for $W \gg N^{3/4}$.

(iv) *Edge statistics.* The transition in eigenvalues statistics is understood at the edge of the spectrum: the prediction from the Thouless criterion was made rigorous by a subtle method of moments. This was proved under the assumption that $\sqrt{2W}(H_{ij})_{i \leq j}$ are ± 1 independent centered Bernoulli random variables, but the method applies to more general distributions. Remember that we order the eigenvalues $\lambda_1 \leq \dots \leq \lambda_N$.

Theorem 3.5 (Transition at the edge of the spectrum [Sodin \[2010\]](#)). *Let $\varepsilon > 0$. If $W \geq N^{\frac{5}{6}+\varepsilon}$, then (2-3) holds. If $W \leq N^{\frac{5}{6}-\varepsilon}$, (2-3) does not hold.*

For eigenvectors (including at the edge of the spectrum), localization cannot hold on less than $W/\log(N)$ entries as proved in [Benaych-Georges and P ech e \[2014\]](#), also by the method of moments.

4 Quantum unique ergodicity and universality


For non mean-field models, eigenvalues and eigenvectors interplay extensively, and their statistics should be understood jointly. Localization (decay of Green's function) is a useful a priori estimate in the proof of Poisson statistics for the Anderson model [Minami \[1996\]](#), and in a similar way we explain below why quantum unique ergodicity implies GOE statistics.

4.1 Mean-field reduction. The method introduced in [Bourgade, Erdős, Yau, and Yin \[2017\]](#) for GOE statistics of band matrices proceeds as follows. We decompose the $1d$ band matrix from (1-1) and its eigenvectors as


$$H = \begin{pmatrix} A & B^* \\ B & D \end{pmatrix}, \quad \mathbf{u}_j := \begin{pmatrix} \mathbf{w}_j \\ \mathbf{p}_j \end{pmatrix},$$

where A is a $W \times W$ matrix. From the eigenvector equation $H\mathbf{u}_j = \lambda_j\mathbf{u}_j$ we have $(A - B^* \frac{1}{D - \lambda_j} B)\mathbf{w}_j = \lambda_j\mathbf{w}_j$. The matrix elements of A do not vanish and thus the above eigenvalue problem features a mean-field random matrix (of smaller size). Hence one can consider the eigenvector equation $Q_e\mathbf{w}_k(e) = \xi_k(e)\mathbf{w}_k(e)$ where

$$(4-1) \quad Q_e = A - B^*(D - e)^{-1}B,$$



(a) A simulation of eigenvalues of $Q_e = A - B^*(D - e)^{-1}B$, i.e. functions $e \mapsto \xi_j(e)$. Here $N = 12$ and $W = 3$. The λ_i 's are the abscissa of the intersections with the diagonal.



(b) Zoom into the framed region of Figure (a), for large N, W : the curves ξ_j are almost parallel, with slope about $1 - N/W$. The eigenvalues of $A - B^*(D - e)^{-1}B$ and those of H are related by a projection to the diagonal followed by a projection to the horizontal axis.

Figure 2: The idea of mean-field reduction: universality of gaps between eigenvalues for fixed e implies universality on the diagonal through parallel projection. For e fixed, we label the curves by $\xi_k(e)$.

and $\lambda_k(e), \mathbf{w}_k(e)$ are eigenvalues and normalized eigenvectors. As illustrated below, the slopes of the functions $e \mapsto \lambda_k(e)$ seem to be locally equal and concentrated:

$$\frac{d}{de} \lambda_k(e) \approx 1 - \frac{1}{\sum_{\alpha=1}^W w_k(\alpha)^2} (1 + o(1)) \approx 1 - \frac{N}{W},$$

which holds for e close to λ_k . The first equality is a simple perturbation formula², and the second is true provided QUE for \mathbf{u}_k holds, in the sense of Equation (3-3) for example.

The GOE local spectral statistics holds for Q_e in the sense (2-2) (it is a mean-field matrix so results from Landon, Sosoe, and Yau [2016] apply), hence it also holds for H by parallel projection: GOE local spectral statistics follow from QUE.

This reduces the problem to QUE for band matrices, which is proved by the same mean-field reduction strategy: on the one hand, by choosing different overlapping blocks A along the diagonal, QUE for H follows from QUE for Q_e by a simple patching procedure (see section 3.3 for more details); on the other hand, QUE for mean-field models is

²The perturbation formula gives a slightly different equation, replacing \mathbf{w}_k by the eigenvector of a small perturbation of H , but we omit this technicality.

known thanks to a strengthening of the eigenvector moment flow method [Bourgade and Yau \[2017\]](#) and [Bourgade, Huang, and Yau \[2017\]](#), explained below.

4.2 The eigenvector moment flow. Obtaining quantum unique ergodicity from the regularity of [Equation \(2-15\)](#) (the eigenvector moment flow) is easy: $\sqrt{N}\langle \mathbf{q}, u_k \rangle$ has limiting Gaussian moments for any \mathbf{q} , hence the entries of u_k are asymptotically independent Gaussian and the following variant of (3-3) holds for Q_e by Markov's inequality (w_k is rescaled to a unit vector): there exists $\varepsilon > 0$ such that for any deterministic $1 \leq k \leq W$ and $I \subset \llbracket 1, W \rrbracket$, for any $\delta > 0$ we have

$$(4-2) \quad \mathbb{P} \left(\left| \sum_{i \in I} |w_k(\alpha)|^2 - \frac{|I|}{N} \right| \geq \delta \right) \leq N^{-\varepsilon/\delta^2}.$$

The main problem with this approach is that the obtained QUE is weak: one would like to replace the above ε with any large $D > 0$, as in (3-3). For this, it was shown in [Bourgade, Yau, and Yin \[2018\]](#) that much more general observables than (2-14) also satisfy the eigenvector moment flow parabolic [Equation \(2-15\)](#).

These new tractable observables are described as follows (we now switch to matrices of dimension N and spectrum $\lambda_1 \leq \dots \leq \lambda_N$, eigenvectors u_1, \dots, u_N , for comparison with (2-14)). Let $I \subset \llbracket 1, N \rrbracket$ be given, $(\mathbf{q}_\alpha)_{\alpha \in I}$ be any family of fixed vectors, and $C_0 \in \mathbb{R}$. Define

$$p_{ij} = \sum_{\alpha \in I} \langle u_i, \mathbf{q}_\alpha \rangle \langle u_j, \mathbf{q}_\alpha \rangle \quad i \neq j \in \llbracket 1, n \rrbracket,$$


$$p_{ii} = \sum_{\alpha \in I} \langle u_i, \mathbf{q}_\alpha \rangle^2 - C_0, \quad i \in \llbracket 1, n \rrbracket,$$

When the \mathbf{q}_α 's are elements of the canonical basis and $C_0 = |I|/N$, this reduces to

$$p_{ij} = \sum_{\alpha \in I} u_i(\alpha) u_j(\alpha), \quad (i \neq j) \quad p_{ii} = \sum_{\alpha \in I} u_i(\alpha)^2 - \frac{|I|}{N}, \quad i \in \llbracket 1, N \rrbracket,$$

and therefore the p_{ij} 's becomes natural partial overlaps measuring quantum unique ergodicity. For any given configuration η as given before (2-14), consider the set of vertices $\mathcal{V}_\eta = \{(i, a) : 1 \leq i \leq n, 1 \leq a \leq 2\eta_i\}$. Let \mathcal{G}_η be the set of perfect matchings of the complete graph on \mathcal{V}_η , i.e. this is the set of graphs G with vertices V_η and edges $\mathcal{E}(G) \subset \{\{v_1, v_2\} : v_1 \in \mathcal{V}_\eta, v_2 \in \mathcal{V}_\eta, v_1 \neq v_2\}$ being a partition of \mathcal{V}_η . For any given edge $e = \{(i_1, a_1), (i_2, a_2)\}$, we define $p(e) = p_{i_1, i_2}$, $P(G) = \prod_{e \in \mathcal{E}(G)} p(e)$ and

$$(4-3) \quad \tilde{f}_{\lambda, t}(\eta) = \frac{1}{\mathfrak{m}(\eta)} \mathbb{E} \left(\sum_{G \in \mathcal{G}_\eta} P(G) \mid \lambda \right), \quad \mathfrak{m}(\eta) = \prod_{i=1}^n (2\eta_i)!!,$$



(a) A configuration η with $\mathfrak{n}(\eta) = 6$, $\eta_{i_1} = 2, \eta_{i_2} = 3, \eta_{i_3} = 1$.

(b) A perfect matching $G \in \mathfrak{G}_\eta$. Here, $P(G) = p_{i_1 i_1} p_{i_1 i_2} p_{i_2 i_2}^2 p_{i_2 i_3} p_{i_3 i_3}$.

where $(2m)!! = \prod_{k \leq 2m, k \text{ odd}} k$. The following lemma is a key combinatorial fact.

Lemma 4.1. *The above function \tilde{f} satisfies the eigenvector moment flow Equation (2-15).*

This new class of observables (4-3) widely generalizes (2-14) and directly encodes the L^2 mass of eigenvectors, contrary to (2-14). Together with the above lemma, one can derive a new strong estimate, (2-14) with some small $\varepsilon > 0$ replaced by any $D > 0$. The mean-field reduction strategy can now be applied in an efficient way: union bounds are costless thanks to the new small N^{-D} error term.

For $d = 2, 3$, the described mean-field reduction together with the strong version of the eigenvector moment flow should apply to give delocalization in some polynomial regime, such as $W \gg N^{99/100}$. However, this is far from the conjectures from (1-2). To approach these transitions, one needs to take more into account the geometry of \mathbb{Z}^d .

4.3 Quantum unique ergodicity and the Gaussian free field. At the heuristic level, the QUE method suggests the transition values W_c from (1-2). More precisely, consider a given eigenvector $u = u_k$ associated to a bulk eigenvalue λ_k . For notational convenience, assume the model’s band width is $2W$ instead of W .

For $\mathbf{1} = (1, \dots, 1) \in \mathbb{Z}^d$, define $\mathcal{W} = \llbracket 1, N \rrbracket^d \cap (2W\mathbb{Z}^d + W\mathbf{1})$. For any $w \in \mathcal{W}$, let $\mathcal{C}_w = \{\alpha \in \mathbb{Z}^d : \|w - \alpha\|_\infty \leq W\}$ be the cell of side length $2W$ around w .

Let $X_w = \sum_{\alpha \in \mathcal{C}_w} u(\alpha)^2$. Consider a set $I, |I| = 2^d$, such that the cells $(\mathcal{C}_w)_{w \in I}$ form an hypercube \mathcal{H} of size $(4W)^d$. Assume one can apply the strong QUE statement (3-3) to a Schur complement Q_e of type (4-1) where A is now chosen to be the $(4W)^d \times (4W)^d$ mean-field matrix indexed by the vertices from \mathcal{H} . We would obtain, for any two adjacent cells $\mathcal{C}_w, \mathcal{C}_v$ with $w, v \in I$,

$$(4-4) \quad \sum_{\alpha \in \mathcal{C}_w} u(\alpha)^2 = \sum_{\alpha \in \mathcal{C}_v} u(\alpha)^2 + \mathcal{O} \left(N^\varepsilon \frac{W^{d/2}}{N^d} \right)$$

with overwhelming probability. By patching these estimates over successive adjacent cells, this gives

$$\sum_{\alpha \in \mathcal{C}_w} u(\alpha)^2 = \left(\frac{W}{N}\right)^d + O\left(N^\varepsilon \frac{W^{d/2}}{N^d}\right) \times \left(\frac{N}{W}\right)^d,$$

and therefore the leading order of $\sum_{\alpha \in \mathcal{C}_w} u(\alpha)^2$ is identified (i.e. QUE holds) for $W \gg N^{\frac{2}{3}}$. This criterion, independent of the dimension d , is more restrictive than (1-2) and omits the important fact that the error term in (4-4) has a random sign.

One may assume that such error terms are asymptotically jointly Gaussian and independent for different pairs of adjacent cells (or at least for such pairs sufficiently far away). We consider the graph with vertices \mathcal{W} and edges the set of pairs (v, w) such that \mathcal{C}_v and \mathcal{C}_w are adjacent cells. A good model for $(X_w)_{w \in \mathcal{W}}$ therefore is a Gaussian vector such that the increments $X_v - X_w$ are independent, with distribution $\mathcal{N}(0, W^d/N^{2d})$ when (v, w) is an edge, and conditioned to (1) $\sum (X_{v_{i+1}} - X_{v_i}) = 0$ for any closed path $v_1, v_2, \dots, v_j, v_1$ in the graph, (2) $X_{v_0} = (W/N)^d$ to fix the ambiguity about definition of X modulo a constant. This model is simply the Gaussian free field, with density for $(X_v)_v$ proportional to

$$e^{-\frac{N^{2d}}{2W^d} \sum_{v \sim w} (x_v - x_w)^2}.$$

As is well known, the Gaussian free field $(Y_v)_v$ on $\llbracket 1, n \rrbracket^d$ with density $e^{-\frac{1}{2} \sum_{v \sim w} (y_v - y_w)^2}$ conditioned to $Y_{v_0} = 0$ has the following typical fluctuation scale, for any deterministic v chosen at macroscopic distance from v_0 (see e.g. [Biskup \[2017\]](#)):

$$(4-5) \quad \text{Var}(Y_v)^{1/2} \approx \begin{cases} n^{1/2} & \text{for } d = 1, \\ (\log n)^{1/2} & \text{for } d = 2, \\ O(1) & \text{for } d = 3. \end{cases}$$

We expect that quantum unique ergodicity (and GOE statistics by the mean-field reduction) holds when $\text{Var}(X_v)^{1/2} \ll \mathbb{E}(X_v)$. With $n = N/W$, this means

$$\frac{W^{d/2}}{N^d} \text{Var}(Y_v)^{1/2} \ll \frac{W^d}{N^d}$$

i.e. $W \gg N^{1/2}$ for $d = 1$, $(\log N)^{1/2}$ for $d = 2$, $O(1)$ for $d = 3$.

Acknowledgments. The author’s knowledge on this topic comes from collaborations with Laszlo Erdős, Horng-Tzer Yau, and Jun Yin. This note reports on joint progress with these authors.

References

- E. Abraham, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan (1979). “Scaling theory of localization: absence of quantum diffusion in two dimensions”. *Phys. Rev. Lett.* 42, p. 673 (cit. on p. 2787).
- M. Aizenman and S. Molchanov (1993). “Localization at large disorder and at extreme energies: an elementary derivation”. *Commun. Math. Phys.* 157, pp. 245–278 (cit. on p. 2787).
- O. Ajanki, L. Erdős, and T. Kruger (2017). “Universality for general Wigner-type matrices”. *Probability Theory and Related Fields* 169 (3–4), pp. 667–727 (cit. on p. 2786).
- (2018). “Stability of the matrix Dyson equation and random matrices with correlations”. *Probability Theory and Related Fields*, pp. 1–81 (cit. on p. 2786).
- N. Anantharaman and E. Le Masson (2015). “Quantum ergodicity on large regular graphs”. *Duke Math. J.* 164.4, pp. 723–765 (cit. on p. 2781).
- G. W. Anderson, A. Guionnet, and O. Zeitouni (2010). *An introduction to random matrices*. Vol. 118. Cambridge Studies in Advanced Mathematics. Cambridge University Press, pp. xiv+492 (cit. on p. 2785).
- G. W. Anderson and O. Zeitouni (2006). “A CLT for a band matrix model”. *Probab. Theory Related Fields* 134.2, pp. 283–338 (cit. on p. 2789).
- P. Anderson (1958). “Absences of diffusion in certain random lattices”. *Phys. Rev.* Pp. 1492–1505 (cit. on pp. 2778, 2787).
- G. Ben Arous and P. Bourgade (2011). “Extreme gaps in the eigenvalues of random matrices”. *Annals of Probability* (cit. on p. 2787).
- Z. Bao and L. Erdős (2017). “Delocalization for a class of random block band matrices”. *Probab. Theory Related Fields* 167.3-4, pp. 673–776 (cit. on p. 2790).
- Z. Bao, L. Erdős, and K. Schnelli (2017). “Local law of addition of random matrices on optimal scale”. *Comm. Math. Phys.* 349.3, pp. 947–990 (cit. on p. 2786).
- R. Bauerschmidt, J. Huang, A. Knowles, and H.-T. Yau (2017). “Bulk eigenvalue statistics for random regular graphs”. *Ann. Probab.* 45.6A, pp. 3626–3663 (cit. on p. 2786).
- R. Bauerschmidt, J. Huang, and H.-T. Yau (2016). “Local Kesten-McKay law for random regular graphs”. *prepublication* (cit. on p. 2792).
- F. Bekeran, A. Figalli, and A. Guionnet (2015). “Transport maps for β -matrix models and universality”. *Comm. Math. Phys.* 338.2, pp. 589–619 (cit. on p. 2787).
- F. Benaych-Georges and S. Péché (2014). “Largest eigenvalues and eigenvectors of band or sparse random matrices”. *Electron. Commun. Probab.* 19, no. 4, 9 (cit. on p. 2793).
- L. Benigni (2017). “Eigenvectors distribution and quantum unique ergodicity for deformed Wigner matrices”. *prepublication* (cit. on p. 2786).
- M. Biskup (2017). “Extrema of the two-dimensional Discrete Gaussian Free Field” (cit. on p. 2797).

- P. Bleher and A. Its (1999). “Semiclassical asymptotics of orthogonal polynomials, Riemann-Hilbert problem, and universality in the matrix model”. *Ann. of Math.* 150, pp. 185–266 (cit. on p. 2786).
- A. Bloemendal, L. Erdős, A. Knowles, H.-T. Yau, and J. Yin (2014). “Isotropic local laws for sample covariance and generalized Wigner matrices”. *Elect. J. Probab.* 19 (cit. on p. 2784).
- L. V. Bogachev, S. A. Molchanov, and L. A. Pastur (1991). “On the density of states of random band matrices”. Russian. *Mat. Zametki* 50.6, pp. 31–42, 157 (cit. on p. 2789).
- P. Bourgade (2018). “Extreme gaps between eigenvalues of Wigner matrices”. *prepublication* (cit. on p. 2787).
- P. Bourgade, L. Erdős, and H.-T. Yau (2014a). “Edge universality for beta ensembles”. *Communications in Mathematical Physics* 332.1, pp. 261–353 (cit. on p. 2786).
- (2014b). “Universality of general β -ensembles”. *Duke Math. J.* 163.6, pp. 1127–1190 (cit. on p. 2786).
- P. Bourgade, L. Erdős, H.-T. Yau, and J. Yin (2016). “Fixed energy universality for generalized Wigner matrices”. *Comm. Pure Appl. Math.* 69.10, pp. 1815–1881 (cit. on pp. 2782, 2783).
- (2017). “Universality for a class of random band matrices”. *Advances in Theoretical and Mathematical Physics* 21 (3), pp. 739–800 (cit. on pp. 2792, 2793).
- P. Bourgade, J. Huang, and H.-T. Yau (2017). “Eigenvector statistics of sparse random matrices”. *Electron. J. Probab.* 22, Paper No. 64, 38 (cit. on pp. 2786, 2795).
- P. Bourgade and H.-T. Yau (2017). “The eigenvector moment flow and local quantum unique ergodicity”. *Comm. Math. Phys.* 350.1, pp. 231–278 (cit. on pp. 2783–2785, 2795).
- P. Bourgade, H.-T. Yau, and J. Yin (2018). “Random band matrices in the delocalized phase I: Quantum unique ergodicity and universality”. *prepublication* (cit. on pp. 2792, 2795).
- S. Brooks and E. Lindenstrauss (2013). “Non-localization of eigenfunctions on large regular graphs”. *Israel J. Math.* 193.1, pp. 1–14 (cit. on p. 2781).
- M.-F. Bru (1989). “Diffusions of perturbed principal component analysis”. *J. Multivariate Anal.* 29.1, pp. 127–136 (cit. on p. 2785).
- G. Casati, L. Molinari, and F. Izrailev (1990Apr). “Scaling properties of band random matrices”. *Phys. Rev. Lett.* 64, pp. 1851–1854 (cit. on p. 2788).
- Z. Che and B. Landon (2017). “Local spectral statistics of the addition of random matrices”. *prepublication* (cit. on p. 2786).
- Y. Colin de Verdière (1985). “Ergodicité et fonctions propres du laplacien”. French, with English summary. *Comm. Math. Phys.* 102.3, pp. 497–502 (cit. on p. 2781).

- P. Deift (1999). *Orthogonal polynomials and random matrices: a Riemann–Hilbert approach*. Vol. 3. Courant Lecture Notes in Mathematics. New York University Courant Institute of Mathematical Sciences, pp. viii+273 (cit. on p. 2786).
- (2017). “Some open problems in random matrix theory and the theory of integrable systems. II”. *SIGMA Symmetry Integrability Geom. Methods Appl.* 13, Paper No. 016, 23 (cit. on p. 2778).
- P. Deift and D. Gioev (2009). *Random matrix theory: invariant ensembles and universality*. Vol. 18. Courant Lecture Notes in Mathematics. Courant Institute of Mathematical Sciences, New York; Amer. Math. Soc., Providence, RI, pp. x+217 (cit. on p. 2786).
- M. Disertori and M. Lager (2017). “Density of states for random band matrices in two dimensions”. *Ann. Henri Poincaré* 18.7, pp. 2367–2413 (cit. on p. 2790).
- M. Disertori, L. Pinson, and T. Spencer (2002). “Density of states for random band matrices”. *Commun. Math. Phys.* 232, pp. 83–124 (cit. on p. 2790).
- F. Dyson (1962). “A Brownian-motion model for the eigenvalues of a random matrix”. *J. Math. Phys.* 3, pp. 1191–1198 (cit. on p. 2783).
- K. Efetov (1997). “Supersymmetry in disorder and chaos”. *Cambridge University Press* (cit. on pp. 2789, 2790).
- L. Erdős and A. Knowles (2011). “Quantum Diffusion and Delocalization for Band Matrices with General Distribution”. *Ann. Inst. H. Poincaré* 12.7, pp. 1227–1319 (cit. on p. 2792).
- (2015). “The Altshuler-Shklovskii formulas for random band matrices I: the unimodular case”. *Comm. Math. Phys.* 333.3, pp. 1365–1416 (cit. on p. 2789).
- L. Erdős, A. Knowles, H.-T. Yau, and J. Yin (2012). “Spectral statistics of Erdős–Rényi graphs II: eigenvalue spacing and the extreme eigenvalues”. *Comm. Math. Phys.* 314, pp. 587–640 (cit. on p. 2786).
- (2013). “Delocalization and diffusion profile for random band matrices”. *Comm. Math. Phys.* 323.1, pp. 367–416 (cit. on pp. 2792, 2793).
- L. Erdős, T. Kruger, and D. Schroder (2017). “Random Matrices with Slow Correlation Decay”. *prepublication* (cit. on p. 2786).
- L. Erdős, S. Péché, J. A. Ramírez, B. Schlein, and H.-T. Yau (2010). “Bulk universality for Wigner matrices”. *Comm. Pure Appl. Math.* 63.7, pp. 895–925 (cit. on p. 2783).
- L. Erdős, B. Schlein, and H.-T. Yau (2009). “Local semicircle law and complete delocalization for Wigner random matrices”. *Commun. Math. Phys.* 287, pp. 641–655 (cit. on p. 2784).
- (2011). “Universality of random matrices and local relaxation flow”. *Inv. Math.* 187.1, pp. 75–119 (cit. on pp. 2782–2784).
- L. Erdős and H.-T. Yau (2015). “Gap universality of generalized Wigner and beta ensembles”. *J. Eur. Math. Soc.* 17, pp. 1927–2036 (cit. on p. 2783).

- L. Erdős, H.-T. Yau, and J. Yin (2012a). “Bulk universality for generalized Wigner matrices”. *Probab. Theory Related Fields* 154.1-2, pp. 341–407 (cit. on pp. [2783](#), [2791](#)).
- (2012b). “Rigidity of eigenvalues of generalized Wigner matrices”. *Adv. Math.* 229.3, pp. 1435–1515 (cit. on pp. [2782](#), [2784](#)).
- D. Facoetti, P. Vivo, and G. Biroli (2016). “From non-ergodic eigenvectors to local resolvent statistics and back: A random matrix perspective”. *EPL (Europhysics Letters)* 115.4 (cit. on p. [2786](#)).
- A. Figalli and A. Guionnet (2016). “Universality in several-matrix models via approximate transportmaps”. *Acta Math.* 217.1, pp. 81–176 (cit. on p. [2787](#)).
- J. Fröhlich and T. Spencer (1983). “Absence of diffusion in the Anderson tight binding model for large disorder or low energy”. *Commun. Math. Phys.* 88, pp. 151–184 (cit. on pp. [2778](#), [2787](#)).
- Y.V. Fyodorov and A.D. Mirlin (1991). “Scaling properties of localization in random bandmatrices: a sigma-model approach”. *Phys. Rev. Lett.* 67 (cit. on pp. [2788](#), [2789](#)).
- I. Goldsheid, S. A. Molchanov, and L. Pastur (1977). “A random homogeneous Schrödinger operator has a pure point spectrum”. *Funkcional. Anal. i Priložen.* 11 (cit. on p. [2787](#)).
- A. Guionnet (2002). “Large deviations upper bounds and central limit theorems for non-commutative functionals of Gaussian large random matrices”. *Annales de l’Institut Henri Poincaré (B)* 38 (3), pp. 341–384 (cit. on p. [2789](#)).
- R. Holowinsky (2010). “Sieving for mass equidistribution”. *Ann. of Math. (2)* 172.2, pp. 1499–1516 (cit. on p. [2781](#)).
- R. Holowinsky and K. Soundararajan (2010). “Mass equidistribution for Hecke eigenforms”. *Ann. of Math. (2)* 172.2, pp. 1517–1528 (cit. on p. [2781](#)).
- J. Huang, B. Landon, and H.-T. Yau (2017). “Transition from Tracy-Widom to Gaussian fluctuations of extremal eigenvalues of sparse Erdős-Rényi graphs”. *prepublication* (cit. on p. [2786](#)).
- I. Jana, K. Saha, and A. Soshnikov (2016). “Fluctuations of linear eigenvalue statistics of random bandmatrices”. *Theory Probab. Appl.* 60.3, pp. 407–443 (cit. on p. [2789](#)).
- K. Johansson (2001). “Universality of the local spacing distribution in certain ensembles of Hermitian Wigner matrices”. *Comm. Math. Phys.* 215.3, pp. 683–705 (cit. on pp. [2782](#), [2783](#)).
- A. Klein (1994). “Absolutely continuous spectrum in the Anderson model on the Bethe lattice”. *Math. Res. Lett.* 1.4, pp. 399–407 (cit. on p. [2788](#)).
- A. Knowles and J. Yin (2013a). “Eigenvector distribution of Wigner matrices”. *Probab. Theory Related Fields* 155.3-4, pp. 543–582 (cit. on p. [2784](#)).
- (2013b). “The isotropic semicircle law and deformation of Wigner matrices”. *Comm. Pure Appl. Math.* 66, pp. 1663–1750 (cit. on p. [2784](#)).

- M. Krishnapur, B. Rider, and B. Virág (2016). “Universality of the stochastic Airy operator”. *Comm. Pure Appl. Math.* 69.1, pp. 145–199 (cit. on p. [2787](#)).
- B. Landon, J. Huang, and H.-T. Yau (2015). “Bulk Universality of Sparse Random Matrices”. *J. Math. Phys.* 56.12 (cit. on p. [2786](#)).
- B. Landon, P. Sosoe, and H.-T. Yau (2016). “Fixed energy universality for Dyson Brownian motion”. *prepublication* (cit. on pp. [2783](#), [2786](#), [2794](#)).
- J.-O. Lee and K. Schnelli (2015). “Local law and Tracy-Widom limit for sparse random matrices”. *to appear in Probab. Theory Related Fields* (cit. on p. [2786](#)).
- J.-O. Lee, K. Schnelli, B. Stetler, and H.-T. Yau (2015). “Bulk universality for deformed Wigner matrices”. *to appear in Annals of Probability* (cit. on p. [2786](#)).
- L. Li and A. Soshnikov (2013). “Central limit theorem for linear statistics of eigenvalues of bandrandom matrices”. *Random Matrices Theory Appl.* 2.4, pp. 1350009, 50 (cit. on p. [2789](#)).
- E. Lindenstrauss (2006). “Invariant measures and arithmetic quantum unique ergodicity”. *Ann. of Math. (2)* 163.1, pp. 165–219 (cit. on p. [2781](#)).
- D. S. Lubinsky (2009). “A new approach to universality limits involving orthogonal polynomials”. *Ann. of Math. (2)* 170.2, pp. 915–939 (cit. on p. [2786](#)).
- M. L. Mehta and M. Gaudin (1960). “On the density of Eigenvalues of a random matrix”. *Nuclear Physics* 18, pp. 420–427 (cit. on p. [2780](#)).
- N. Minami (1996). “Local fluctuation of the spectrum of a multidimensional Anderson tightbinding model”. *Commun. Math. Phys.* 177, pp. 709–725 (cit. on pp. [2778](#), [2788](#), [2793](#)).
- J. R. Norris, L. C. G. Rogers, and David Williams (1986). “Brownian motions of ellipsoids”. *Trans. Amer. Math. Soc.* 294.2, pp. 757–765 (cit. on p. [2785](#)).
- L. Pastur and M. Shcherbina (1997). “Universality of the local eigenvalue statistics for a class of unitary invariant random matrix ensembles”. *J. Stat. Phys.* 86.1-2, pp. 109–147 (cit. on p. [2786](#)).
- R. Peled, J. Schenker, M. Shamis, and S. Sodin (2017). “On the Wegner orbital model”. *International Mathematical Research Notices* (cit. on p. [2790](#)).
- Z. Rudnick and P. Sarnak (1994). “The behaviour of eigenstates of arithmetic hyperbolic manifolds”. *Comm. Math. Phys.* 161.1, pp. 195–213 (cit. on p. [2781](#)).
- J. Schenker (2009). “Eigenvector localization for random band matrices with power law-band width”. *Comm. Math. Phys.* 290, pp. 1065–1097 (cit. on pp. [2789](#), [2790](#)).
- M. Shcherbina (2014). “Change of variables as a method to study general β -models: Bulk universality”. *J. Math. Phys.* 55 (cit. on p. [2787](#)).
- M. Shcherbina and T. Shcherbina (2017). “Characteristic polynomials for 1D random band matrices from the localization side”. *Communications in Mathematical Physics* 351 (cit. on p. [2790](#)).

- (2018). “Universality for 1d Random Band Matrices: Sigma-Model Approximation”. *Journal of Statistical Physics*, pp. 1–38 (cit. on p. 2791).
- T. Shcherbina (2014a). “On the Second Mixed Moment of the Characteristic Polynomials of 1D Band Matrices”. *Communications in Mathematical Physics* 328, pp. 45–82 (cit. on pp. 2790, 2791).
- (2014b). “Universality of the local regime for the block band matrices with a finite number of blocks”. *J. Stat. Phys.* 155.3, pp. 466–499 (cit. on p. 2790).
- A. I. Shnirel’man (1974). *Uspekhi Mat. Nauk* 29.6, pp. 181–182 (cit. on p. 2781).
- S. Sodin (2010). “The spectral edge of some random band matrices”. *Ann. of Math.* 173.3, pp. 2223–2251 (cit. on pp. 2789, 2793).
- (2014). “Several applications of the moment method in random matrix theory”. *Proceedings of the International Congress of Mathematicians* (cit. on p. 2789).
- P. von Soosten and S. Warzel (2017). “Non-Ergodic Delocalization in the Rosenzweig-Porter Model”. *prepublication* (cit. on p. 2786).
- T. Spencer (2010). “Random banded and sparse matrices (Chapter 23) in “Oxford Handbook of Random Matrix Theory” edited by G. Akemann, J. Baik, and P. Di Francesco” (cit. on p. 2789).
- (2012). “SUSY statistical mechanics and random band matrices” (cit. on pp. 2788–2790).
- T. Tao and V. Vu (2011). “Random matrices: universality of local eigenvalue statistics”. *Acta Math.* 206.1, pp. 127–204 (cit. on pp. 2782–2784).
- (2012). “Random matrices: universal properties of eigenvectors”. *Random Matrices Theory Appl.* 1.1 (cit. on p. 2784).
- D. J. Thouless (1977). “Maximum metallic resistance in thin wires”. *Physical Review Letters* 39.18, pp. 1167–1169 (cit. on p. 2789).
- C. Tracy and H. Widom (1994). “Level Spacing Distributions and the Airy Kernel”. *Communications in Mathematical Physics* 159, pp. 151–174 (cit. on p. 2780).
- H. F. Trotter (1984). “Eigenvalue distributions of large Hermitian matrices; Wigner’s semi-circle law and a theorem of Kac, Murdock, and Szegő”. *Adv. in Math.* 54.1, pp. 67–82 (cit. on p. 2778).
- W.-M. Wang (1992). *On localization and density of states for the random Schroedinger-operator*. Thesis (Ph.D.)—Princeton University. ProQuest LLC, Ann Arbor, MI, p. 92 (cit. on p. 2789).
- E. Wigner (1957). “Distribution of neutron resonance level spacing”. *In International conference on the neutron interactions with the nucleus (Columbia University, New York, 1957), Columbia Univ. Rept. CU-175 (TID-7547)*, pp. 49–50 (cit. on p. 2778).
- S. Zelditch (1987). “Uniform distribution of eigenfunctions on compact hyperbolic surfaces”. *Duke Math. J.* 55.4, pp. 919–941 (cit. on p. 2781).

Received 2018-03-05.

PAUL BOURGADE
COURANT INSTITUTE
NEW YORK UNIVERSITY
bourgade@cims.nyu.edu