


# The curve and the Langlands program

Laurent Fargues (CNRS/IMJ)



## The curve

- ▶ Defined and studied in our joint work with Fontaine

# The curve

- ▶ Defined and studied in our joint work with Fontaine
- ▶ Two aspects : compact  $p$ -adic Riemann surface / algebraic curve

# The curve

- ▶ Defined and studied in our joint work with Fontaine
- ▶ Two aspects : compact  $p$ -adic Riemann surface / algebraic curve
- ▶ Starting datum :  $F|\mathbb{F}_p$  perfectoid field

## The curve

- ▶ Defined and studied in our joint work with Fontaine
- ▶ Two aspects : compact  $p$ -adic **Riemann surface** / **algebraic curve**
- ▶ Starting datum :  $F|\mathbb{F}_p$  perfectoid field
- ▶  $p$ =variable,  $Y = \{0 < |p| < 1\}$  open punctured disk, adic space

$$W(\mathcal{O}_F)\left[\frac{1}{p}, \frac{1}{[\varpi]}\right] = \left\{ \sum_{n \gg -\infty} [x_n] p^n \mid x_n \in F, \sup_n |x_n| < +\infty \right\}$$

## The curve

- ▶ Defined and studied in our joint work with Fontaine
- ▶ Two aspects : compact  $p$ -adic **Riemann surface** / **algebraic curve**
- ▶ Starting datum :  $F|\mathbb{F}_p$  perfectoid field
- ▶  $p$ =variable,  $Y = \{0 < |p| < 1\}$  open punctured disk, adic space

$$W(\mathcal{O}_F)\left[\frac{1}{p}, \frac{1}{[\varpi]}\right] = \left\{ \sum_{n \gg -\infty} [x_n] p^n \mid x_n \in F, \sup_n |x_n| < +\infty \right\}$$

- ▶  $\mathcal{O}(Y)$  completion of the preceding. Fontaine's ring. No explicit formula for elements of  $\mathcal{O}(Y)$

# The curve

►  $Y \xrightarrow{\varphi}$  Frobenius

$$X = Y/\varphi^{\mathbb{Z}}$$

compact adic space  $/\mathbb{Q}_p$

# The curve

- ▶  $Y \xrightarrow{\varphi}$  Frobenius

$$X = Y/\varphi^{\mathbb{Z}}$$

compact adic space  $/\mathbb{Q}_p$

- ▶  $\mathcal{O}(1)$  line bundle  $/X \leftrightarrow$  automorphic factor  $j(\varphi, p) = p$



# The curve

- ▶  $Y \xrightarrow{\varphi}$  Frobenius

$$X = Y/\varphi^{\mathbb{Z}}$$

compact adic space  $/\mathbb{Q}_p$

- ▶  $\mathcal{O}(1)$  line bundle  $/X \leftrightarrow$  automorphic factor  $j(\varphi, p) = p$
- ▶ Schematical curve

$$\mathfrak{X} = \text{Proj}(\oplus_{d \geq 0} \mathcal{O}(Y)^{\varphi = p^d})$$

# The curve

- ▶  $Y \xrightarrow{\varphi}$  Frobenius

$$X = Y/\varphi^{\mathbb{Z}}$$

compact adic space  $/\mathbb{Q}_p$

- ▶  $\mathcal{O}(1)$  line bundle  $/X \leftrightarrow$  automorphic factor  $j(\varphi, p) = p$
- ▶ Schematical curve

$$\mathfrak{X} = \text{Proj}\left(\bigoplus_{d \geq 0} \mathcal{O}(Y)^{\varphi = p^d}\right)$$

- ▶ GAGA type morphism

$$X \longrightarrow \mathfrak{X}$$

# The curve

## Theorem (F.-Fontaine)

- ▶ *The curve is a curve* :  $\mathfrak{X}$  is a Dedekind scheme,  $X$  noetherian regular dimension 1 adic space

# The curve

## Theorem (F.-Fontaine)

- ▶ *The curve is a curve* :  $\mathfrak{X}$  is a Dedekind scheme,  $X$  noetherian regular dimension 1 adic space
- ▶ *Perfectoid residue fields* :

$$\forall x \in |\mathfrak{X}|, k(x)|\mathbb{Q}_p \text{ perfectoid, } [k(x)^{\flat} : F] < +\infty$$

# The curve

## Theorem (F.-Fontaine)

- ▶ *The curve is a curve* :  $\mathfrak{X}$  is a Dedekind scheme,  $X$  noetherian regular dimension 1 adic space
- ▶ *Perfectoid residue fields* :

$$\forall x \in |\mathfrak{X}|, k(x) | \mathbb{Q}_p \text{ perfectoid, } [k(x)^{\flat} : F] < +\infty$$

- ▶  $\widehat{\mathcal{O}}_{\mathfrak{X},x} = B_{dR}^+(k(x))$

# The curve

## Theorem (F.-Fontaine)

- ▶ *The curve is a curve* :  $\mathfrak{X}$  is a Dedekind scheme,  $X$  noetherian regular dimension 1 adic space
- ▶ *Perfectoid residue fields* :

$$\forall x \in |\mathfrak{X}|, k(x) | \mathbb{Q}_p \text{ perfectoid, } [k(x)^b : F] < +\infty$$

- ▶  $\widehat{\mathcal{O}}_{\mathfrak{X},x} = B_{dR}^+(k(x))$
- ▶  $\mathfrak{X}$  is "complete" :  $\deg(x) := [k(x)^b : F]$

$$\forall f \in \mathbb{Q}_p(\mathfrak{X})^\times, \deg(\operatorname{div} f) = 0$$

## The curve

- ▶  $\mathfrak{X} \setminus \{x\} = \text{Spec}(B_x)$ ,  $B_x$  Dedekind, P.I.D. if  $F$  alg. closed.  
 $(B_x, -ord_x)$  **not euclidean** i.e.

$$H^1(\mathcal{O}(-1)) \neq 0$$

## The curve

- ▶  $\mathfrak{X} \setminus \{x\} = \text{Spec}(B_x)$ ,  $B_x$  Dedekind, P.I.D. if  $F$  alg. closed.  
 $(B_x, -ord_x)$  **not euclidean** i.e.

$$H^1(\mathcal{O}(-1)) \neq 0$$

- ▶  $F$  alg. closed.  $\Rightarrow |\mathfrak{X}| = \text{untlts of } F \text{ up to Frob.}$



## Vector bundles on the curve

- ▶ GAGA verified :  $\text{Bun}_{\mathfrak{X}} \xrightarrow{\sim} \text{Bun}_X$

## Vector bundles on the curve

- ▶ GAGA verified :  $\text{Bun}_{\mathfrak{X}} \xrightarrow{\sim} \text{Bun}_X$
- ▶  $\mathfrak{X}$  complete  $\Rightarrow$  good Harder-Narasimhan reduction theory

$$\mu = \frac{\text{deg}}{\text{rk}}$$

## Vector bundles on the curve

- ▶ GAGA verified :  $\text{Bun}_{\mathfrak{X}} \xrightarrow{\sim} \text{Bun}_X$
- ▶  $\mathfrak{X}$  complete  $\Rightarrow$  good Harder-Narasimhan reduction theory

$$\mu = \frac{\text{deg}}{\text{rk}}$$

- ▶  $\lambda \in \mathbb{Q} \rightsquigarrow \mathcal{O}(\lambda)$  stable slope  $\lambda$  vector bundle,  $\lambda = \frac{d}{h}$ ,  
pushforward of  $\mathcal{O}(d)$  via cyclic cover  $Y/\varphi^{h\mathbb{Z}} \rightarrow Y/\varphi^{\mathbb{Z}}$

## Vector bundles on the curve

- ▶ GAGA verified :  $\text{Bun}_{\mathfrak{X}} \xrightarrow{\sim} \text{Bun}_X$
- ▶  $\mathfrak{X}$  complete  $\Rightarrow$  good Harder-Narasimhan reduction theory

$$\mu = \frac{\text{deg}}{\text{rk}}$$

- ▶  $\lambda \in \mathbb{Q} \rightsquigarrow \mathcal{O}(\lambda)$  stable slope  $\lambda$  vector bundle,  $\lambda = \frac{d}{h}$ ,  
pushforward of  $\mathcal{O}(d)$  via cyclic cover  $Y/\varphi^{h\mathbb{Z}} \rightarrow Y/\varphi^{\mathbb{Z}}$

### Theorem (F.-Fontaine)

*F alg. closed*

$$\begin{aligned} \{\lambda_1 \geq \dots \geq \lambda_n \mid \lambda_i \in \mathbb{Q}\} &\xrightarrow{\sim} \text{Bun}_{\mathfrak{X}} / \sim \\ (\lambda_1, \dots, \lambda_n) &\mapsto [\oplus_i \mathcal{O}(\lambda_i)]. \end{aligned}$$

## Vector bundles on the curve : applications

- ▶ Quick simple proofs of the two fundamental theorems of  $p$ -adic Hodge theory :
  - ▶ weakly admissible  $\Rightarrow$  admissible (Colmez-Fontaine)
  - ▶ de Rham  $\Rightarrow$  pot. semi-stable (Berger)

## Vector bundles on the curve : applications

- ▶ Quick simple proofs of the two fundamental theorems of  $p$ -adic Hodge theory :
  - ▶ weakly admissible  $\Rightarrow$  admissible (Colmez-Fontaine)
  - ▶ de Rham  $\Rightarrow$  pot. semi-stable (Berger)
- ▶  $F = \mathbb{C}_p^b$ ,

$$\mathfrak{X} \curvearrowright \text{Gal}(\overline{\mathbb{Q}}_p | \mathbb{Q}_p)$$

$\infty \in |\mathfrak{X}|$  fixed. Study **Galois equivariant vector bundles** and their **modifications** at  $\infty$

# Vector bundles on the curve : applications

- ▶ Quick simple proofs of the two fundamental theorems of  $p$ -adic Hodge theory :
  - ▶ weakly admissible  $\Rightarrow$  admissible (Colmez-Fontaine)
  - ▶ de Rham  $\Rightarrow$  pot. semi-stable (Berger)
- ▶  $F = \mathbb{C}_p^b$ ,

$$\mathfrak{X} \curvearrowright \text{Gal}(\overline{\mathbb{Q}}_p | \mathbb{Q}_p)$$

$\infty \in |\mathfrak{X}|$  fixed. Study Galois equivariant vector bundles and their modifications at  $\infty$

- ▶ Look at modifications of vector bundles
  - ▶  $\varphi$ -modules over  $A_{inf}$
  - ▶ Scholze-Weinstein

## $G$ -bundles

- ▶  $G$  reductive group /  $\mathbb{Q}_p$ ,  $\check{\mathbb{Q}}_p := \widehat{\mathbb{Q}_p^{un}}$ ,  $\sigma = \text{Frob}$ ,

$$B(G) = G(\check{\mathbb{Q}}_p) / \sigma\text{-conj}, \quad b \sim gbg^{-\sigma}$$



## G-bundles

- ▶  $G$  reductive group /  $\mathbb{Q}_p$ ,  $\check{\mathbb{Q}}_p := \widehat{\mathbb{Q}_p^{un}}$ ,  $\sigma = \text{Frob}$ ,

$$B(G) = G(\check{\mathbb{Q}}_p) / \sigma\text{-conj}, \quad b \sim gbg^{-\sigma}$$

- ▶  $b \in G(\check{\mathbb{Q}}_p)$

$$\mathcal{E}_b = Y \times_{\varphi} G$$

$G$ -bundle /  $X$ ,  $\varphi$  acts on  $G$  via conjugation by  $b\sigma$

## G-bundles

- ▶  $G$  reductive group /  $\mathbb{Q}_p$ ,  $\check{\mathbb{Q}}_p := \widehat{\mathbb{Q}_p^{un}}$ ,  $\sigma = \text{Frob}$ ,  
 $B(G) = G(\check{\mathbb{Q}}_p) / \sigma\text{-conj}$ ,  $b \sim gbg^{-\sigma}$

- ▶  $b \in G(\check{\mathbb{Q}}_p)$

$$\mathcal{E}_b = Y \times_{\varphi} G$$

$G$ -bundle /  $X$ ,  $\varphi$  acts on  $G$  via conjugation by  $b\sigma$

### Theorem (F)

$F$  alg. closed

$$\begin{aligned} B(G) &\xrightarrow{\sim} H_{\text{ét}}^1(\mathcal{X}, G) \\ [b] &\mapsto [\mathcal{E}_b] \end{aligned}$$

## G-bundles

- ▶  $G$  reductive group /  $\mathbb{Q}_p$ ,  $\check{\mathbb{Q}}_p := \widehat{\mathbb{Q}_p^{un}}$ ,  $\sigma = \text{Frob}$ ,  
 $B(G) = G(\check{\mathbb{Q}}_p) / \sigma\text{-conj}$ ,  $b \sim gbg^{-\sigma}$

- ▶  $b \in G(\check{\mathbb{Q}}_p)$

$$\mathcal{E}_b = Y \times_{\varphi} G$$

$G$ -bundle /  $X$ ,  $\varphi$  acts on  $G$  via conjugation by  $b\sigma$

### Theorem (F)

$F$  alg. closed

$$\begin{aligned} B(G) &\xrightarrow{\sim} H_{\text{ét}}^1(\mathcal{X}, G) \\ [b] &\mapsto [\mathcal{E}_b] \end{aligned}$$

- ▶ Dictionary : **reduction theory** (Atiyah-Bott) for  $G$ -bundles / Kottwitz description of  $B(G)$ .
- ▶ *Example* :  $\mathcal{E}_b$  semi-stable  $\Leftrightarrow b$  is basic (isoclinic)

## $G$ -bundles : applications

- ▶ (Chen, F, Shen) : **Proof of F.-Rapoport conjecture on period spaces** ( $p$ -adic analogs of Griffith's period spaces).  
Example : computation of  $p$ -adic period spaces for  $SO(2, n - 2)$

## $G$ -bundles : applications

- ▶ (Chen, F, Shen) : **Proof of F.-Rapoport conjecture on period spaces** ( $p$ -adic analogs of Griffith's period spaces).  
Example : computation of  $p$ -adic period spaces for  $SO(2, n - 2)$
- ▶ **Construction of local Shimura varieties** (Scholze)  $\text{Sht}(G, b, \mu)$   
as moduli of modifications  $\mathcal{E}_1 \rightsquigarrow \mathcal{E}_b$

## $G$ -bundles : applications

- ▶ (Chen, F, Shen) : **Proof of F.-Rapoport conjecture on period spaces** ( $p$ -adic analogs of Griffith's period spaces).  
Example : computation of  $p$ -adic period spaces for  $SO(2, n - 2)$
- ▶ **Construction of local Shimura varieties** (Scholze)  $\mathrm{Sht}(G, b, \mu)$   
as moduli of modifications  $\mathcal{E}_1 \rightsquigarrow \mathcal{E}_b$
- ▶ **Newton stratification of Hodge-Tate flag manifold**  
(Caraiani-Scholze), modification  $\mathcal{E}_{1,x}$ ,  $x \in$  flag manifold,  
 $\mathcal{E}_{1,x} \simeq \mathcal{E}_b$  for some  $b \rightsquigarrow$  stratification by the set of such  $[b]$

# The stack $\text{Bun}_G$ of $G$ -bundles/curve

- ▶ Introduced to formulate a **geometrization conjecture of the local Langlands correspondence**

$$\underbrace{\varphi}_{\text{Langlands parameter}} \quad \longmapsto \quad \underbrace{\mathcal{F}_\varphi}_{\text{perverse sheaf on } \text{Bun}_G}$$

# The stack $\text{Bun}_G$ of $G$ -bundles/curve

- ▶ Introduced to formulate a **geometrization conjecture of the local Langlands correspondence**

$$\underbrace{\varphi}_{\text{Langlands parameter}} \quad \longmapsto \quad \underbrace{\mathcal{F}_\varphi}_{\text{perverse sheaf on } \text{Bun}_G}$$

- ▶  $S = \overline{\mathbb{F}}_p$ -perfectoid space  $\rightsquigarrow X_S$  adic space/ $\mathbb{Q}_p =$  "family of curves  $(X_{k(s)})_{s \in S}$ "



# The stack $\mathrm{Bun}_G$ of $G$ -bundles/curve

- ▶ Introduced to formulate a **geometrization conjecture of the local Langlands correspondence**

$$\underbrace{\varphi}_{\text{Langlands parameter}} \quad \longmapsto \quad \underbrace{\mathcal{F}_\varphi}_{\text{perverse sheaf on } \mathrm{Bun}_G}$$

- ▶  $S = \overline{\mathbb{F}}_p$ -perfectoid space  $\rightsquigarrow X_S$  adic space/ $\mathbb{Q}_p =$  "family of curves  $(X_{k(s)})_{s \in S}$ "
- ▶  $\mathrm{Bun}_G = v$ -stack on  $\mathrm{Perf}_{\overline{\mathbb{F}}_p}$  of  $G$ -bundles/curve

$$B(G) \xrightarrow{\sim} |\mathrm{Bun}_G|$$

$$\mathrm{Bun}_G = \coprod_{\alpha \in \pi_1(G)_\Gamma} \mathrm{Bun}_G^{c_1 = \alpha}$$

# The stack $\text{Bun}_G$



$$\text{Bun}_G^{c_1=0,ss} = [\bullet / \underline{G(\mathbb{Q}_p)}] \xrightarrow{\text{open}} \text{Bun}_G$$

# The stack $\text{Bun}_G$



$$\text{Bun}_G^{c_1=0,ss} = [\bullet / \underline{G(\mathbb{Q}_p)}] \xrightarrow{\text{open}} \text{Bun}_G$$

- ▶  $\underline{\text{Aut}}(\text{trivial } G\text{-bundle}) = \underline{G(\mathbb{Q}_p)}$  topological group  $\neq$  classical case  $\rightsquigarrow G$  algebraic group

# The stack $\text{Bun}_G$



$$\text{Bun}_G^{c_1=0,ss} = [\bullet / \underline{G(\mathbb{Q}_p)}] \xrightarrow{\text{open}} \text{Bun}_G$$

▶  $\underline{\text{Aut}}(\text{trivial } G\text{-bundle}) = \underline{G(\mathbb{Q}_p)}$  topological group  $\neq$  classical case  $\rightsquigarrow G$  algebraic group

▶ More generally  $\forall \alpha \in \pi_1(G)_\Gamma$

$$\text{Bun}_G^{c_1=\alpha,ss} = [\bullet / \underline{J_b(\mathbb{Q}_p)}]$$

$b$  basic,  $J_b$  inner form of  $G$

# The stack $\text{Bun}_G$



$$\text{Bun}_G^{c_1=0,ss} = [\bullet / \underline{G(\mathbb{Q}_p)}] \xrightarrow{\text{open}} \text{Bun}_G$$

- ▶  $\underline{\text{Aut}}(\text{trivial } G\text{-bundle}) = \underline{G(\mathbb{Q}_p)}$  topological group  $\neq$  classical case  $\rightsquigarrow G$  algebraic group

- ▶ More generally  $\forall \alpha \in \pi_1(G)_\Gamma$

$$\text{Bun}_G^{c_1=\alpha,ss} = [\bullet / \underline{J_b(\mathbb{Q}_p)}]$$

$b$  basic,  $J_b$  inner form of  $G$

- ▶ In general each H.N. stratum is a classifying stack

$$[\bullet / \mathcal{J}_b]$$

$$\mathcal{J}_b = \mathcal{J}_b^0 \times \underline{J_b(\mathbb{Q}_p)}, \mathcal{J}_b^0 = \text{unipotent diamond}$$

# The geometrization conjecture

- ▶  $\varphi =$  discrete Langlands parameter

$$\varphi : W_{\mathbb{Q}_p} \longrightarrow {}^L G$$

# The geometrization conjecture

- ▶  $\varphi =$  discrete Langlands parameter

$$\varphi : W_{\mathbb{Q}_p} \longrightarrow {}^L G$$

- ▶ Conjecture

$$\varphi \longmapsto \mathcal{F}_\varphi$$

$S_\varphi$ -equivariant perverse Hecke eigensheaf on  $\text{Bun}_G$

- ▶ s.t. the stalks of  $\mathcal{F}_\varphi$  at semi-stable points gives local Langlands + internal structure of L-packets for all extended pure inner forms of  $G$

# The geometrization conjecture

For this :

- ▶ Need to give a meaning to "perverse sheaf on  $\text{Bun}_G$ "
- ▶ Need to give a meaning to the Hecke eigensheaf property : establish *geometric Satake* in this context



# The geometrization conjecture

For this :

- ▶ Need to give a meaning to "perverse sheaf on  $\text{Bun}_G$ "
- ▶ Need to give a meaning to the Hecke eigensheaf property : establish **geometric Satake** in this context

↪ **joint work with Scholze** : give a precise statement of the conjecture + construction of the local Langlands correspondence

$$\pi \mapsto \varphi_\pi$$

à la V. Lafforgue using  $\text{Bun}_G$

# Constructible and perverse étale sheaves on $\text{Bun}_G$

Joint with Scholze.

$\Lambda \in \{\overline{\mathbb{F}}_\ell, \overline{\mathbb{Q}}_\ell\}$  étale coefficients,  $\ell \neq p$

# Constructible and perverse étale sheaves on $\mathrm{Bun}_G$

Joint with Scholze.

$\Lambda \in \{\overline{\mathbb{F}}_\ell, \overline{\mathbb{Q}}_\ell\}$  étale coefficients,  $\ell \neq p$

- ▶  $\mathrm{Bun}_G$  is a (cohomologically) **smooth stack of dimension 0**

# Constructible and perverse étale sheaves on $\text{Bun}_G$

Joint with Scholze.

$\Lambda \in \{\overline{\mathbb{F}}_\ell, \overline{\mathbb{Q}}_\ell\}$  étale coefficients,  $\ell \neq p$

- ▶  $\text{Bun}_G$  is a (cohomologically) **smooth stack of dimension 0**
- ▶ Good notion of **constructible sheaf** on  $\text{Bun}_G =$  reflexive sheaves w.r.t. Verdier duality

# Constructible and perverse étale sheaves on $\text{Bun}_G$

Joint with Scholze.

$\Lambda \in \{\overline{\mathbb{F}}_\ell, \overline{\mathbb{Q}}_\ell\}$  étale coefficients,  $\ell \neq p$

- ▶  $\text{Bun}_G$  is a (cohomologically) **smooth stack of dimension 0**
- ▶ Good notion of **constructible sheaf** on  $\text{Bun}_G =$  reflexive sheaves w.r.t. Verdier duality
- ▶ Fiberwise criterion of constructibility in terms of representation theory :  $\forall b \in G(\check{\mathbb{Q}}_p)$ , the stalk at the point given by  $b$  is an **admissible** representation of  $J_b(\mathbb{Q}_p)$

# Geometric Satake

- ▶  $\mathrm{Div}^1 = \mathrm{Spa}(\mathbb{Q}_p)^\diamond / \varphi^{\mathbb{Z}}$  sheaf of deg. 1 effective div. on the curve

# Geometric Satake

- ▶  $\text{Div}^1 = \text{Spa}(\mathbb{Q}_p)^\diamond / \varphi^{\mathbb{Z}}$  sheaf of deg. 1 effective div. on the curve



$$\text{Gr}_G^{B_{dR}} \longrightarrow \text{Spa}(\mathbb{Q}_p)^\diamond$$

Scholze's  $B_{dR}$  affine grassmanian,

$$\text{Gr}_G^{B_{dR}} / \varphi^{\mathbb{Z}} \rightarrow \text{Div}^1$$

Beilinson-Drinfeld type affine Grassmanian

# Geometric Satake

- ▶  $\mathrm{Div}^1 = \mathrm{Spa}(\mathbb{Q}_p)^\diamond / \varphi^{\mathbb{Z}}$  sheaf of deg. 1 effective div. on the curve



$$\mathrm{Gr}_G^{B_{dR}} \longrightarrow \mathrm{Spa}(\mathbb{Q}_p)^\diamond$$

Scholze's  $B_{dR}$  affine grassmanian,

$$\mathrm{Gr}_G^{B_{dR}} / \varphi^{\mathbb{Z}} \rightarrow \mathrm{Div}^1$$

Beilinson-Drinfeld type affine Grassmanian

Theorem (F.-Scholze)

*Geometric Satake holds for  $\mathrm{Gr}_G^{B_{dR}}$ , Satake category  $\simeq \mathrm{Rep}({}^L G, \Lambda)$ .*



# Construction of Langlands parameters

- ▶ Factorization enhancement :

$$I \text{ finite set} \rightsquigarrow \mathrm{Gr}_{G,I}^{B_{dR}} \rightarrow (\mathrm{Spa}(\mathbb{Q}_p)^\diamond)^I$$

+ factorization property when  $I$  varies.

# Construction of Langlands parameters

- ▶ Factorization enhancement :

$$I \text{ finite set} \rightsquigarrow \mathrm{Gr}_{G,I}^{B_{dR}} \rightarrow (\mathrm{Spa}(\mathbb{Q}_p)^\diamond)^I$$

+ factorization property when  $I$  varies.



$$\begin{array}{ccc} & \mathrm{Hecke}_I & \\ & \swarrow \quad \searrow & \\ \mathrm{Bun}_G & & \mathrm{Bun}_G \times (\mathrm{Spa}(\mathbb{Q}_p)^\diamond)^I \end{array}$$

# Construction of Langlands parameters

- ▶ Factorization enhancement :

$$I \text{ finite set} \rightsquigarrow \mathrm{Gr}_{G,I}^{B_{dR}} \rightarrow (\mathrm{Spa}(\mathbb{Q}_p)^\diamond)^I$$

+ factorization property when  $I$  varies.



$$\begin{array}{ccc} & \mathrm{Hecke}_I & \\ & \swarrow \quad \searrow & \\ \mathrm{Bun}_G & & \mathrm{Bun}_G \times (\mathrm{Spa}(\mathbb{Q}_p)^\diamond)^I \end{array}$$

- ▶  $W \in \mathrm{Rep}({}^L G^I, \Lambda) \rightsquigarrow IC_W$  kernel on  $\mathrm{Hecke}_I$  via geo Satake

# Construction of Langlands parameters

- ▶ Coupled with V. Lafforgue strategy (global function field) we construct local Langlands

$$\underbrace{\pi}_{\text{irred. rep. of } G(\mathbb{Q}_p)} \longmapsto \underbrace{\varphi_\pi}_{\text{semi-simple Langlands parameter}}$$

# Construction of Langlands parameters

- ▶ Coupled with V. Lafforgue strategy (global function field) we construct local Langlands

$$\underbrace{\pi}_{\substack{\text{irred. rep.} \\ \text{of } G(\mathbb{Q}_p)}} \longmapsto \underbrace{\varphi_\pi}_{\substack{\text{semi-simple Langlands} \\ \text{parameter}}}$$

- ▶ Very little known about this correspondence : surjectivity, finiteness of fibers ?

# Construction of Langlands parameters

- ▶ Coupled with V. Lafforgue strategy (global function field) we construct local Langlands

$$\underbrace{\pi}_{\substack{\text{irred. rep.} \\ \text{of } G(\mathbb{Q}_p)}} \longmapsto \underbrace{\varphi_\pi}_{\substack{\text{semi-simple Langlands} \\ \text{parameter}}}$$

- ▶ Very little known about this correspondence : surjectivity, finiteness of fibers ?
- ▶ Geometrization conjecture goes in the other direction

Langlands parameter  $\longmapsto$  representation

and would give this + **internal structure of L-packets**

## Back to the geometrization conjecture

- ▶ The  $GL_1$ -case. Classically : Abel-Jacobi morphism **locally trivial fibration in simply connected alg. var.** (projective spaces) in high degree

## Back to the geometrization conjecture

- ▶ The  $GL_1$ -case. Classically : Abel-Jacobi morphism *locally trivial fibration in simply connected alg. var.* (projective spaces) in high degree
- ▶ Reduced to the following theorem

### Theorem (F)

For  $d \geq 3$ , the Abel-Jacobi morphism

$$AJ^d : \text{Div}^d \longrightarrow \mathcal{P}ic^d$$

is a *pro-étale locally trivial fibration in simply connected diamonds*.

Here

$$\text{Div}^d = \text{Hilbert diamond} = (\text{Div}^1)^d / \mathfrak{S}_d$$

with  $\text{Div}^1 = \text{Spa}(\mathbb{Q}_p)^\diamond / \varphi^{\mathbb{Z}}$ .



That's only the beginning of the story!